

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/57, C07K 14/47, C12N 9/64, A61K 38/48, 38/17		A1	(11) International Publication Number: WO 97/32981 (43) International Publication Date: 12 September 1997 (12.09.97)
(21) International Application Number: PCT/GB97/00603 (22) International Filing Date: 4 March 1997 (04.03.97) (30) Priority Data: 9604865.7 7 March 1996 (07.03.96) GB 9611896.3 7 June 1996 (07.06.96) GB 9614293.0 8 July 1996 (08.07.96) GB 9624028.8 19 November 1996 (19.11.96) GB (71) Applicant (for all designated States except US): IMUTRAN LIMITED [GB/GB]; Douglas House, 18 Trumpington Road, Cambridge CB2 2AH (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): FARRIES, Timothy, Charles [GB/GB]; The MRC Centre, Hills Road, Cambridge CB2 2QH (GB). HARRISON, Richard, Alexander [GB/GB]; The MRC Centre, Hills Road, Cambridge CB2 2QH (GB). (74) Agents: WHITE, Martin, Paul et al.; Kilburn & Strode, 30 John Street, London WC1N 2DD (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: DOWN-REGULATION RESISTANT C3 CONVERTASE			
(57) Abstract Native complement pathway proteins modified such that the protein is capable of forming a down-regulation resistant C3 convertase. Preferably the modified protein is a modified human C3 protein. DNA sequences encoding such proteins are also provided, together with DNA constructs. Conjugates comprising such proteins and a specific binding moiety, for example an antibody, are also described, as are uses of such proteins and/or conjugates in therapy.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DOWN-REGULATION RESISTANT C3 CONVERTASE

5 The present invention relates to novel modified proteins capable of forming C3 convertases resistant to down-regulation, DNA sequences encoding such proteins and the use of such proteins as therapeutic agents, particularly for use in depleting levels of complement pathway proteins or in targeting complement attack (C3b deposition) at specific sites.

10

The complement system functions in the immune response of humans and other vertebrates, being of major importance in the effector functions such as phagocytosis, cytolysis and recruitment of cells that induce local inflammatory responses [15]. These properties are desirable for elimination of invading pathogens, such as bacteria, but undesirable when triggered to act against host tissues (e.g. in post-ischemic reperfusion injury [3]) or against foreign therapeutic material (e.g. hyperacute rejection of xenografts [7]). There have been attempts to abrogate these undesirable properties by exploiting derivatives of complement regulatory proteins whose normal function is to suppress complement activation [10, 18].

20

25 The complement system comprises proteins both on the surface of cells, (receptors and regulators) as well as in the fluid-phase (blood plasma and other extracellular environments). The critical step for the generation of responses is the proteolytic conversion of C3 to the fragments C3b and C3a. C3a is an anaphylatoxin that, like C5a, attracts mast cells to the site of challenge, resulting in local release of histamine, vasodilation and other inflammatory effects. The nascent C3b has an ability to bind to surfaces around its site of

30

generation. This C3b then focuses attack by the cytolytic complement components (C5-C9).

5 Surface-bound C3b, and its degradation products, also function as ligands for C3 receptors mediating, for example, phagocytosis [15]. There are two distinct pathways of complement activation that both result in conversion of C3 to C3b and subsequent responses. The classical pathway is commonly triggered by complexes of antibody with antigen, initiating an enzyme cascade
10 involving the proteins C1q, C1r, C1s, C2 and C4. The alternative pathway depends on an activation loop involving C3 itself and requiring factors B and D.

15 Conversion of C3 to C3b (or C3i) produces a product that can combine with factor B, giving C3bB (or C3iB). These complexes are acted upon by factor D to generate C3bBb, which is a C3 convertase capable of cleaving more C3 to C3b, leading to more C3bBb and even more C3 conversion.
20 Under certain circumstances the C3bBb complex is stabilised by association with the positive regulator properdin (P). However, this positive-feedback loop is normally limited to a slow tick-over by regulatory proteins, notably factor H and factor I.

25 Factor H (and structurally related cell-associated molecules) (i) displaces B and Bb from C3b, and (ii) acts as a cofactor for factor I which cleaves C3b into iC3b thereby preventing any recombination with factor B to form more C3 convertases. The pathway is "fired" into
30 amplified generation of C3b in the presence of surfaces, such as many bacterial cell walls, that bind nascent C3b and impede its regulation by factors H and I. Nascent C3b is also able to bind to endogenous cells. Endogenous cell

surfaces normally exposed to complement are therefore additionally protected by membrane-bound regulators such as MCP, DAF and CR1 acting in a similar manner to factor H.

5

There are a few rare naturally occurring conditions where the normal fluid-phase regulation cannot occur and spontaneous C3 conversion ultimately results in generalised depletion of C3 from the circulation:- (i) genetic deficiencies of factor H or I [13], (ii) the presence of antibodies (nephritic factors) that bind to C3bBb and impede dissociation [4], and (iii) contact with a protein in cobra venom, called cobra venom factor (CVF), that combines with factor B and forms a C3 convertase enzyme which does not contain C3b and is not affected by factors H and I [14]. These illustrate the normal physiological importance of down-regulation of complement in the absence of specific activation.

There are also circumstances where specific activation occurs, but is unwanted, particularly when it is directed against tissues of the host (e.g. tissue damaged by ischemia or surgery) or against foreign material deliberately given for therapeutic purposes (such as a xenograft, artificial organ or a dialysis membrane). The complement activation results in undesirable attack and further damage, so in these cases it would be beneficial to block or inhibit the activation and response.

Existing approaches to preventing complement-mediated damage have targeted the use of down-regulatory proteins (CR1, MCP, DAF and factors H and I) to inhibit complement activation. Complement inhibitors like factor I, factor H and soluble derivatives of the membrane-bound proteins

CR1, DAF, MCP do suppress the fluid-phase amplification loop of the alternative pathway. Therefore there have been attempts to use these molecules, particularly CR1 (which seems to be the most potent) to reduce complement-mediated damage in models of physiological situations [10, 18].

Factor H is endogenously present in blood plasma in high concentrations (typically 0.3-0.5 mg/ml [15]), so even though increased levels of inhibitors do dampen-down fluid-phase reactions, their potency is weak so large amounts of purified proteins would have to be administered in vivo (e.g. probably in excess of 5 mg/Kg body weight of soluble CR1). In addition, the alternative pathway is activated by surfaces where the effect of factor H is already impeded. While this does not necessarily concomitantly reduce the activities of other inhibitors, the same factors suggest that they are unlikely to be completely or universally effective.

Cobra Venom Factor (CVF) has the property of generating a stable C3 convertase which can be used experimentally to deplete complement in animals in vivo, and in other samples (e.g. human blood plasma) in vitro. CVF is potent (e.g. 40 µg/Kg can destroy the complement activity of a mouse [16]). However, there are disadvantages that make it unsuitable for therapeutic use in humans.

Firstly, it is obtained from cobra venom (a difficult source to obtain and dangerous to handle) and must therefore be carefully purified from the venom neurotoxins. There is also the obvious difficulty in obtaining supplies. This problem cannot readily be overcome by cloning and expressing the gene ex vivo,

because there are post-translational modifications that occur in the snake (specific proteolytic processing) that may be difficult (or impossible) to reproduce *in vitro*. In addition, the enzymes and digestion conditions required for this processing are currently unknown. Secondly, the protein is of foreign origin (to humans) and therefore immunogenic. This precludes its repeated therapeutic use, as would be required to deplete a patient over many weeks (e.g. to allow xenograft survival).

Although CVF has some structural and functional homologies with human C3 [17], it also has major differences in both respects (e.g. chain structure, site of biosynthesis, insensitivity to complement regulators, formation of a stable C3 convertase). It is not derived from the cobra equivalent of C3 which is known, having been cloned and sequenced, and which in gross structure and function resembles human C3 more closely than does CVF [8].

CVF is a venom-specific product of an animal of great evolutionary distance from *homo sapiens*. It is therefore not practicable to use genetic manipulation to modify this protein into a product that can be used non-immunogenically in humans.

We have now devised an alternative strategy which relies on by-passing the physiological regulation and, instead of inhibiting complement activation, causes the system to be super-activated. This has two applications. Firstly, it can be used *in vivo* to activate complement until one or more components are exhausted, resulting in loss of ability to produce local responses to any subsequent

challenge (such as a xenograft). Secondly, the unregulated super-activation can be deliberately localised to a particular target (e.g. a virus or a virally-infected cell) to increase the sensitivity of that target to complement-mediated destructive responses.

The term "regulators of complement activation" is used herein to include all proteins that act to inhibit amplification of C3 conversion, and is not intended to be restricted in meaning to those proteins whose genes are located in the RCA genetic locus. It does not however include "up-regulators" such as properdin. "C3 conversion" is defined as the proteolytic conversion of C3 into C3b and C3a, unless otherwise indicated, and "C3 convertase" (or simply "convertase") is defined as an enzyme (typically a complex of two or more protein components; for example C3bBb, C3iBb, CVFBb or C4b2a) that catalyses this reaction.

Thus, in a first aspect the invention provides a native complement pathway protein modified such that the protein is capable of forming a down-regulation resistant C3 convertase.

By "native" is meant naturally occurring, ie is obtainable in nature. Thus, the definition encompasses any naturally occurring complement pathway protein modified as defined above. It is not intended to be restricted to species specific proteins. In other words, a modified human protein could be used as a down-regulation resistant C3 convertase in other mammalian species, for example. Typically, modified complement pathway proteins from the same species will be used.

Modification of the C3 DNA coding sequence, for example using site directed mutagenesis, can produce a variant of C3 that is resistant to complement regulatory proteins while retaining positive functional properties (cleavage to C3b by C3 convertase) and features of structural integrity (correct chain structure, and presence of a thiolester bond). The invention described herein relates to genetically-modified forms of native complement proteins, for example human C3, whose C3b fragment acquires the property of being resistant to physiological complement regulation. Because of this resistance, these molecules can generate stabilised forms of the corresponding C3 convertase that produce amplified conversion of C3 to C3b, and later degradation products, in physiological environments (e.g. in vivo).

In a preferred embodiment the invention provides a modified human C3 protein which is resistant to cleavage by factor I.

This can be achieved by modifying residues of the protein at proteolytic sites.

A particularly preferred embodiment of the invention relates to a modified human C3 protein wherein the protein is modified by replacement of either Arg-1303, Arg-1320 or both by another amino acid. The other amino acid may be Tyrosine, Cystine, Tryptophan, Glutamine, Glutamic acid or Glycine. Arg-1303 is preferably replaced by Glutamic acid or Glycine (less preferably by Glutamine). Arg-1320 is preferably replaced by Glutamine.

Other strategies for producing suitable modified proteins

of the invention include:

5 i) Reduced susceptibility to the inhibitory actions of factor H and related proteins (eg. MCP, DAF, CR1). For example, in human C3 residues 767-776 and 1209-1271 have been implicated in factor H binding [20,24], and substitution of one or more of these residues or other residues also associated with the action of these proteins, could reduce the binding of one or more of
10 these regulatory proteins.

ii) Reduced rate of dissociation of C3bBb. Mutations can be introduced which would strengthen the interaction between C3b and Bb. This would result in both a reduction
15 in spontaneous decomposition of the enzyme, and diminish the effectiveness of factor H (and related regulators) in displacing Bb from C3b.

These mutations are desirable to reduce the rates of both
20 the spontaneous and the factor H-mediated decomposition of C3bBb. Even in the absence of factor H, the fluid phase C3bBb complex has a half-life of only about 10 mins at 37°C in the presence of properdin [6].

25 iii) Human C3 residues 752-761 are implicated in binding factor B. It is a highly conserved region in C3, and a closely related sequence is found in C4. As C4 binds the factor B homolog C2, the strong similarity of this region between C3 and C4, together with its high conservation in
30 C3, further supports its role in C3 as a factor B binding site. Thus, changes in this region could have effects on B affinity and on the stability of C3bBb.

iv) Resistance to other regulators of complement

activation such as CR1, DAF and MCP would also be desirable. The mode of action of these regulators are all similar to factor H, so additional mutagenesis would not necessarily be required. Similarly, some pathogenic organisms express their own inhibitors of complement activation that are often structurally and functionally homologous to factor H (e.g. *Vaccinia* virus secretory peptide []). These molecules protect the invaders against immune responses, and it would be advantageous to be able to attack them with targeted C3 convertase enzymes resistant to these defences.

v) Mutations that increase the stabilisation of the C3 convertase by properdin. The activity of properdin is to stabilise the C3bBb complex, retarding spontaneous and factor H-dependent dissociation. This stabilisation is ineffective in the fluid-phase, but seems to be more important in amplifying the process once it has already started on a suitable activating surface [5]. Increasing its activity (by increasing its affinity) may upset the balance in the fluid-phase, and thereby promote spontaneous C3 conversion. This should be particularly useful in combination with the other modifications described above.

vi) Mutations that prevent the C3bBb from possessing C5 convertase activity. When used to deplete active C3 from the circulation an undesirable side-effect could be the generation of large amounts of anaphylactic peptides. The most potent of these is C5a, which is cleaved from C5 by some C3 convertase enzymes. This reaction probably depends on the affinity of the convertase for another molecule of C3b [11], and so may be subject to suppression by mutations to the C3 that remove this

interaction.

vii) Improved activity of the C3 convertase. The active site of the C3bBb C3 convertase enzyme resides in the Bb portion. The C3b component presumably functions to impose an active conformation on Bb and/or to bind and orientate the substrate to be acted upon by Bb. This is not known, but in either case there may be scope for enhancing the activity of the convertase through mutations in C3.

viii) Expression in a functional form. Wild-type C3 requires conversion to C3b before it can combine into a new C3 convertase complex. When used in vivo, a requirement for conversion to C3b (or C3i) would delay the action of the modified C3. It would therefore be desirable to either administer the protein in a form capable of immediate convertase formation, or to administer pre-formed convertase complexes. It is therefore advantageous to generate a functionally C3b-like reagent ex-vivo. This could be achieved in vitro (e.g. by proteolysis).

ix) Modifications to the native protein which serve to introduce new cleavage sites such that peptide regions required for factor B binding are retained but those required exclusively for factor H binding can be specifically removed. For example, sites can be introduced such that the C3b-like form of the modified C3 can be further cleaved into a form that still binds factor B but is less susceptible to inactivation by factors H and I.

x) Modifications in other regions which may affect the

C3b interaction with factor B and/or factor H.

5 The invention is based on reversing the traditional approach by promoting C3 conversion to deplete C3 and thereby disable the system. An additional application of the invention is the potential to promote C3 conversion at a particular site, and thereby recruit the complement-dependent effector mechanisms to attack a specific target.

10

Therefore the ultimate effect will be to increase the amount of C3 conversion when the modified protein is administered into a physiological medium (e.g. blood) containing regulators of complement activation. This activity can then be used either to deplete that medium of native C3, or to localise the C3 conversion at a desired target.

15

20 The analogue of C3 whose C3b-fragment is resistant to the actions of factor I (e.g. the derivative described in example 1) would bind factor B, which would then be cleaved by factor D and eventually dissociate in an inactive form. In the absence of inactivation by factor I, the modified C3b would be able to repeatedly bind new molecules of factor B and thereby promote its inactivation. Therefore another potential application of modifications described in this invention would be the inactivation of the alternative pathway by consumption of factor B activity. An analogous approach could also be used to modify C4 to promote the consumption of C2, and thereby disable the classical pathway of complement activation.

25

30

The invention includes any other protease used in an

analogous manner to the C3bBb enzyme which leads to cleavage of C3 to C3b, despite the presence of regulators of complement activation.

5 The invention also includes DNA sequences which code for a protein of the invention as well as DNA constructs comprising such DNA sequences.

10 "DNA sequences" include all other nucleic acid sequences which, by virtue of the degeneracy of genetic code, also code for the given amino acid sequence or which are substantially homologous to this sequence. These sequences are thus also included within the scope of the invention.

15 Nucleic acid sequences which are "substantially homologous" are also within the scope of the present invention. "Substantial homology" may be assessed either at the nucleic acid level or at the amino acid level. At
20 the nucleic acid level, sequences having substantial homology may be regarded as those which hybridise to the nucleic acid sequences of the invention under stringent conditions (for example, at 35 to 65°C in a salt solution of about 0.9M). At the amino acid level, a protein
25 sequence may be regarded as substantially homologous to another protein sequence if a significant number of the constituent amino acids exhibit homology. At least 55%, 60%, 70%, 80%, 90%, 95% or even 99%, in increasing order of preference, of the amino acids may be homologous.

30 As discussed above the proteins of the invention can be used to achieve localised complement activation effects. One way of ensuring this is to conjugate the protein to a moiety which will bind at the desired target. Thus, in

another aspect the invention provides a conjugate comprising a protein of the invention linked to a specific binding moiety, for example a specific binding protein. An example of such a protein would be an antibody or an antigen binding fragment thereof.

The proteins of the invention are intended to be administered to a subject to elicit a desired therapeutic effect. To that end therefore the invention also provides:

- a) A protein of the invention for use in therapy;
- b) The use of a protein or a conjugate of the invention in the manufacture of a medicament for use in depleting levels of complement pathway protein, and in particular for use in preventing rejection of foreign matter;
- c) A pharmaceutical formulation comprising one or more proteins or conjugates of the invention together with one or more pharmaceutically acceptable carriers and/or excipients; and
- d) A method of reducing complement pathway protein in a mammal which comprises administering to the mammal a protein of the invention, preferably in the form of a pharmaceutical formulation.

Pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per dose. Such a unit may contain as a minimum, for example, 1mg of active ingredient, and preferably 2-3mg. The upper limit which such a unit dose can contain will depend on many factors such as the

condition being treated, the route of administration and the age, weight and condition of the patient as well as economic considerations. As an example a unit dose form can contain as much as 10mg or even 100mg of active ingredient.

The proteins of the invention could be used in vivo to disable the complement system. Circumstances where this may be desirable include the following:-

(a) In order to prevent complement-mediated destruction or damage to a transplant, particularly a xenograft (material transplanted from a different species of animal), and especially a discordant xenograft (where the donor and recipient species are discordantly related). The recipient would be decumented prior to the operation and maintained in this state until the transplant had either been accommodated or been replaced by a more compatible organ.

The initial treatment could be made within several days before transplantation. Additional decumentation could be required at times of rejection crisis. The treatments may be accompanied by the use of anti-histamine reagents to control the general inflammatory responses (e.g. vasodilation) likely to result from the generation of C3a and/or C5a.

Decumentation may also be beneficial in the use of artificial organs or tissues (e.g. artificial kidney dialysis membranes) which activate the complement system. As described above, the protein may be given either as the unactivated form, a functionally C3b-like form or a preformed active C3 convertase (like C3bBb). These may be

administered by any route whereby the active convertase will encounter the circulating C3 (e.g. intravenously, subcutaneously etc.).

5 Another alternative would be an ex vivo treatment, for example by transfusing the circulation through a matrix bearing the active convertase. This could have the advantage of allowing anaphylactic peptides (C3a and C5a) and other low molecular weight inflammatory mediators
10 (e.g. histamine and nitric oxide) to be removed (e.g. by dialysis) prior to the decompartmented blood (or plasma) being returned to the patient.

(b) To prevent complement-mediated damage resulting from
15 major surgery. The patient would be decompartmented, as above, preferably before the operation (but if necessary afterwards) and kept in this state until the danger of additional internal injury due to complement-dependent immune attack had diminished.

20 (c) To minimise complement-mediated damage resulting from non-surgical injury. In these cases the decompartmentation must be performed after the initial injury, but the formulations and methods of
25 administration are likely to be otherwise similar to those described above. This may be particularly useful when the recovery involves reperfusion of an ischemic tissue by the circulation (e.g. myocardial ischemia, frostbite, burns etc.).

30 (d) To minimise complement-mediated damage resulting from antibody-antigen interactions. Complement-mediated defensive responses are particularly undesirable in autoimmune diseases which may include glomerulonephritis,

haemolytic anaemia, myasthenia gravis, Diabetes type I, rheumatoid arthritis and multiple sclerosis. Disabling the complement system during severe episodes of disease may alleviate the condition, for instance by local administration to the joint in rheumatoid arthritis.

(e) To make a specific pathogenic target more susceptible to complement-mediated immune mechanisms. In this approach, the aim is not to use the super-active C3 convertase to produce generalised depletion of C3, but instead to use the convertase locally to concentrate the C3 conversion at a desired target. The target may be a pathogenic organism, such as a bacteria, virus or other parasite, or a deleterious host cell or tissue, such as a tumour cell or a virally-infected cell. The C3 convertase could be localised to the target either by local administration (e.g. direct injection, possibly in a medium that retards its dispersion into the general circulation), or by combining with a targeting moiety, e.g. an antibody. Thus the modified protein could be linked to a specific immunoglobulin either by chemical cross-linking of the proteins, or by joining the DNA coding sequences and expressing (and purifying) the fusion protein (e.g. in the case of IgG, either the heavy or the light chain could be attached to C3 and co-expressed with C3, or both chains could be combined within one complete fusion polypeptide), or by incorporation of specific coding sequences (eg. for "leucine zipper"-like domains) to the DNA of both fusion partners (eg. modified C3 and specific antibody) such that the expressed products, when mixed together, self-associate to form stable conjugates. The fusion protein could then be administered locally or into the general circulation.

Liposomes (bearing the antibody on the surface with the modified protein either on the surface or inside the liposome) and/or virions (e.g. engineered to express the proteins on their surface) could also be used for co-delivery of antibody and modified protein. This strategy could be used directly, alone or in combination with other treatments, at any stage in the disease process. It may be particularly appropriate for use in eliminating any cancerous cells left in the circulation after surgical removal of a tumour. The antibody-modified protein conjugates could also be used *ex vivo* to eliminate pathogenic tissue. For example to kill leukaemic cells from an extracted bone-marrow and then returning the remaining healthy cells to the patient.

Alternatively lymphocytes that do not match the MHC types of the recipient could be eliminated from a bone marrow prior to transplantation. Also the modified protein could be linked to an antigen, and this combination could be used, either *in vivo* or *ex vivo*, to attack lymphocytes of undesirable reactivities (e.g. against transplant or self tissue).

The same technology would be applicable to treating other species, using either a human modified protein derivative, or a similar analogue tailor-made for that species.

Preferred features of each aspect of the invention are as for each other aspect *mutatis mutandis*.

The invention will now be described by way of the following examples, which should not be construed as in any way limiting the invention. The examples refer to the

accompanying drawings in which:

Figure 1: shows the predicted protein sequence of human C3 as encoded in PC3;
(using the standard one letter amino acid code)

Figure 2: shows the cDNA sequence in PC3;
(using the standard one letter deoxynucleotide code for the sense strand, written 5'-3').

Figure 3: shows a visualisation of modified proteins of the invention.

Figure 4: shows the effect of various mutations to human C3 which replace Arg 1303 or Arg 1320 on factor I-mediated cleavage at these sites.

N.B.

1. [35S]-biosynthetically labelled samples.
2. Reactions performed at normal ionic strength.
3. Immunoprecipitated with anti-C3.
4. SDS-PAGE under reducing conditions.
5. Autoradiography.

Figure 5: shows enhanced resistance of human C3 incorporating the Arg 1303 -> Gln 1303 mutation to inactivation by factors I and H.

Figure 6: shows an analysis of the cleavage of a C3 convertase mutated at amino acid residues 752-754 and 758-760.

This is a photograph of a Western Blot developed from a 7.5% polyacrylamide SDS-PAGE gel (reducing conditions), after electrophoretic transfer onto

5 nitrocellulose, probing with a sheep anti-human C3 antibody, and development with horse-radish-peroxidase-coupled anti-sheep Immunoglobulin antibody and Enhance ChemiLuminescence (method and detection reagents from Amersham, U.K.) recorded on X-ray film. The cleavage reactions and detection procedure were performed as described in Example 4 with reference to the results shown in Figure 3.

10

Key:

Tracks 1-4: wild-type C3 (expressed in COS cells)
Tracks 5-8: Mutant C3 (residues 752-754 changed to Gly-Ser-Gly and residues 758-760 also being changed to Gly-Ser-Gly) (expressed in COS cells)

15

Tracks 1,5: no addition
Tracks 2,6: + CVFBb
Tracks 3,7: + factors H + I
Tracks 4,8: + CVFBb + factors H + I

20

The bands indicated by arrows are:

A: C3 alpha-chain
B: C3 alpha'-chain
C: C3 beta chain
25 D: 68 kDa cleavage product of C3 alpha'-chain
E: IgG heavy chain

30

Figure 7: shows an analysis of the cleavage of radiolabelled factor B by factor D, in the presence of wild-type and mutant C3's (C3i's)

A photograph of the autoradiograph of the SDS-PAGE gel is shown. All samples contained factor D and ¹²⁵I-labelled factor B, and were incubated for 3

hours at 37°C.

The samples in the numbered tracks also included:

1. Buffer alone
- 5 2. 1/125 wild-type C3
3. 1/25 wild-type C3
4. 1/5 wild-type C3
5. 1/25 mutant C3 (residues 1427 Gln, 1431 Asp and 1433 Gln)
- 10 6. 1/5 mutant C3
7. undiluted mutant C3

The bands indicated by arrows are:

- A. Uncleaved ^{125}I -labelled factor B (93 kDa)
- 15 B. 60 kDa cleavage product ("Bb")
- C. 33 kDa cleavage product ("Ba")

Figure 8: shows an SDS-PAGE study illustrating the formation of a conjugate between C3i and IgG.

20

This is a Coomassie stain of a 4% acrylamide SDS-PAGE gel run under non-reducing conditions. The numbered tracks contain samples of:

1. PDP-IgG
- 25 2. C3i
3. PDP-IgG + C3i reaction mixture

Indicated by arrows are:

- A. Probably C3i-IgG conjugate (350 kDa)
- 30 B. C3i (200 kDa)
- C. IgG (150 kDa)

Figure 9: demonstrates that conjugate targets C3 convertase activity against sheep erythrocytes.

(This graph shows the % lysed sheep erythrocytes after coating with dilutions of either the C3i-IgG conjugate, PDP-IgG or C3i followed by washing, generation of C3 convertases with properdin and factors B and D, and finally development of lysis by NGPS in CFD/EDTA, as described in the methods. Only the conjugate produces lysis, and this lysis is dose dependent.)

5
10 Figures 10 and 11: show the cleavage properties of the DV-1AM mutant C3 (see Examples 12-14).

15 In respect of Figure 10, COS cell supernatants containing expressed wild-type (A) and DV-1AM mutant (B) C3 were treated with 1) -; 2) CVFBb; 3) 10 µg/ml factor I and 50 µg/ml factor H; or 4) CVFBb plus 10 µg/ml factor I and 50 µg/ml factor H, immunoprecipitated, analysed by SDS-PAGE (in the lanes indicated) and electroblotted onto nitrocellulose as described in example 4. In this case
20 the blot was developed using a combination of rat monoclonal antibodies, Clone-3 and Clone-9, that-react with the C3dg region of C3 and its fragmentation products (Lachmann, P.J. et al, *J Immunol*, 41:503 (1980)), followed by a horse radish peroxidase-coupled anti rat immunoglobulin (from Sigma) and detection using the ECL
25 reagents and procedure supplied by Amersham.

30 In respect of Figure 11, COS cell supernatants containing expressed DV-1B mutant (A), wild-type (B) and DV-6 mutant (C) C3 were treated with 1) -; 2) 10 µg/ml factor I and 50 µg/ml factor H; 3) CVFBb; 4) CVFBb plus 10 µg/ml factor I and 2 µg/ml factor H; 5) CVFBb plus 10 µg/ml factor I and 10 µg/ml factor H; or 6) CVFBb plus 10 µg/ml factor I and 50 µg/ml factor H; immunoprecipitated,

- 22/1 -

analysed by SDS-PAGE (in the lanes indicated), electroblotted onto nitrocellulose and detected using a polyclonal sheep anti-C3 antibody as described in example 4.

Figure 12 shows an analysis of a C3 related product resulting from a frame shift mutation. This product is referred to as HDV-3X. Results for HDV-3X are compared with results for DV-3, which like HDV-3X includes the mutations T1031G, E1032N, Q1033H, E1035N and K1036I, but unlike HDV-3X is not modified at the C-terminus.

COS cell supernatants containing expressed DV-3 (A) and HDV-3X mutant (B) C3 were treated with 1) -; 2) CVFBb + 10µg/ml Factor I; 3) CVFBb + 10µg/ml Factor I + 1 µg/ml Factor H; 4) CVFBb + 10µg/ml Factor I + 5 µg/ml Factor H; 5) CVFBb + 10µg/ml Factor I + 25 µg/ml Factor H; immunoprecipitated, analysed by SDS-PAGE (in the lanes indicated) and electroblotted onto nitrocellulose as described in example 4. In this case the blot was developed using a combination of rat monoclonal antibodies, Clone-3 and Clone-9, that react with the C3dg region of C3 and its fragmentation products (Lachmann, P.J. et al, 1980, J. Immunol. 41:503), followed by a horse radish peroxidase-coupled anti rat immunoglobulin (from Sigma) and detection using the ECL reagents and procedure supplied by Amersham.

Figure 13 is in respect of an experiment where COS cell supernatants containing expressed E1Q2(A), E1Q2QG3(B) and E1Q2E3(C) mutant C3 which were treated (37°C, 2.5 hr) with 1) CVFBb + 10µg/ml Factor I; 2) CVFBb + 10µg/ml Factor I + 50µg/ml Factor H; 3) CVFBb + 10µg/ml Factor I + 25µg/ml sCR1; immunoprecipitated, analysed by SDS-PAGE (in the lanes indicated) and electroblotted onto nitrocellulose as described in example 4. In this case the blot was developed using a combination of rat monoclonal antibodies, Clone-3 and Clone-9, as described in example 12, followed by a horse radish peroxidase-coupled anti rat immunoglobulin (from Sigma) and detection using the ECL reagents and procedure supplied by Amersham.

- 22/2 -

The 86 kDa band (arrowed) is the product of Factor I mediated cleavage at the third site. when there has been no prior cleavage at sites 1 or 2.

Figure 14 is in respect of an experiment where COS cell supernatants containing expressed NC3 (A), FT-1 (B), FT-2 (C), FT-3 (D), FT-4 (E) and FT-5 (F) mutant C3 were treated (37°C, 2.75 hr) with 1) -; 2) CVFBB + 10µg/ml Factor I + 50µg/ml Factor H; immunoprecipitated, analysed by SDS-PAGE (in the lanes indicated) and electroblotted onto nitrocellulose as described in example 4. The blot was developed using a combination of rat monoclonal antibodies, Clone-3 and Clone-9, as described in example 12, followed by a horse radish peroxidase-coupled anti rat immunoglobulin (from Sigma) and detection using the ECL reagents and procedure supplied by Amersham.

Figure 15 is in respect of an experiment where COS cell supernatants containing expressed NC3 (A), FR-1 (B), FR-2 (C), FR-3 (D), FR-4 (E) and FT-2 (F) mutant C3 were treated (37°C, 2.5 hr) with 1) -; 2) CVFBB + 10µg/ml Factor I + 50µg/ml Factor H; immunoprecipitated, analysed by SDS-PAGE (in the lanes indicated) and electroblotted onto nitrocellulose as described in example 4. The blot was developed using a combination of rat monoclonal antibodies, Clone-3 and Clone-9, as described in example 12, followed by a horse radish peroxidase-coupled anti rat immunoglobulin (from Sigma) and detection using the ECL reagents and procedure supplied by Amersham.

- 22/3 -

For ease of reference, the relationship between the claims on file which relate to particular proteins capable of acting as down-regulation resistant C3 convertases and the examples and tables provided hereafter is set out below:

TABLE A

Claims	Region of amino acid sequence (relative to human C3 convertase) believed to be important for down- regulation resistance	Example	Table
5-8	1303/1320	1-6, 11	
9-11	758-780/752-754	7	
12	1427, 1431, 1433	8	I
13-15	992-1005	12, 13	II
16-19	1152-1155	14	II
20-29	1546-1663	15, 17	III
30-34	954-955	16	

The following standard methods and definitions are applicable to all the examples:

All complement components referred to are of human origin, unless otherwise specified, using standard terminology for all proteins and their derived fragments

(e.g. as contained in reference [15])). In addition the term "C3i" refers to any molecular form of C3 without an intact thiolester bond, but retaining the C3a polypeptide on the alpha chain.

The human C3 cDNA and coding sequence are numbered as shown in FIGURE 2, using the numbering used in the EMBL nucleotide data base (derived from reference [2]). The sequence shown is that of our construct ('PC3'), which lacks the first 11 nucleotides of the 5' untranslated region reported in reference [2], and hence the first base is numbered 12. The putative initiation codon is nucleotides number 61-63, the codon for the amino-terminal serine residue of the beta-chain is nucleotides 127-129, and the codon for the amino-terminal serine residue of the alpha-chain is nucleotides 2074-2076.

The protein sequence is numbered according to the precursor sequence as shown in FIGURE 1, which is a predicted translation of the DNA sequence in Appendix 1 (amino acids 1-22 are expected to comprise a signal sequence that is removed during biosynthesis, and amino

acids 668-671 are expected to be removed when the precursor is cleaved into the alpha and beta chains).

The following abbreviations have the following meanings;
5 CVP cobra venom factor; ELISA, Enzyme-linked
immunoassorbant assay; *E. coli*, *Escherichia coli*; kb,
kilobase; HSV-1, herpes simplex virus type 1; PBS,
phosphate-buffered saline. COS-1 is a cell line derived
from monkey kidney cells. The following are restriction
10 endonucleases:- AflIII, DraI, DraIII, EcoRI, EcoRV,
HindIII, NaeI, NheI, XbaI.

Standard methods

15 Methods for standard molecular biological procedures such
as plasmid isolations, agarose gel electrophoresis, and
DNA ligations can be found in reference [21]. Double
stranded DNA was sequenced using the 'Sequenase version
2.0' kit supplied by 'United States Biochemicals'. C3
20 expression was measured by an ELISA assay using plastic
plates pre-coated with affinity-purified polyclonal sheep
anti-human C3 to which samples of culture supernatant
were added. Bound C3 was detected with a monoclonal rat
antibody to C3 conjugated to alkaline phosphatase, and
25 the chromogenic substrate, p-nitrophenol phosphate.
Assays were calibrated with purified human plasma C3.

Methods for purification of complement proteins and CVP,
and for the preparation of affinity purified anti-C3
30 antibodies used in the analysis can be found in reference
[28]. Equivalent reagents can also be purchased from
Sigma chemical company LTD.

C3 cDNA coding sequence

Our C3 cDNA coding sequence was constructed from two segments isolated from a random-primed human liver cDNA library carried in the vector pGEM4 (Promega). Five oligodeoxynucleotides, corresponding to known segments in the human C3 coding sequence, were radiolabeled with T4 polynucleotide kinase and [γ -32P]ATP and used to probe filter transfers of the library from agarose plates. Two clones containing inserts of approximately 4 kb were isolated. Restriction endonuclease digestion, hybridisation to specific oligodeoxynucleotide probes and partial sequence analysis demonstrated that one of these ('A13') included the 5'-end of the 5.1kb message, whereas the other ('B44') extended to the 3'-end.

These inserts therefore overlapped by approximately 3 kb, including a unique *EcoRI* restriction enzyme site. The incomplete 5' section of A13 was cut out with *EcoRI* and *NheI*, and replaced with the complete segment isolated from B44 by digestion with *EcoRI* and *XbaI*. Both pieces were purified by gel electrophoresis in low-melting point agarose before ligating together with T4 DNA ligase to produce a vector ('PGC3') containing 5.1 kb of DNA encoding the entire C3 precursor protein.

Linker sequences 5' to the C3 coding region contained two ATG's which are potential false translation start sites. These were therefore removed by gapped-plasmid mutagenesis, as described in the method of example 1, using an oligodeoxynucleotide PL-ATC-3 (tagggagacc ggaagcttgc cctctccctc tgctccctctg t) that deleted approximately 50 base pairs of linker/adaptor DNA, without altering the C3 coding sequence. This mutated vector, 7.7kb containing 5.1kb of C3 cDNA sequence plus 2.6 kb of sequence from the pGEM4 vector (Promega) is

referred to as PC3.

5 The C3 coding region of the PGC3 plasmid was completely sequenced and revealed only four differences from a previously published human C3 ("S" allele) cDNA sequence [2].

(i) the changes C2481->G, and C2805->T do not alter the coding;

10 (ii) T1001->C encodes the previously described HAV 4-1- (Leucine314->Proline) polymorphic form [20]; and

(iii) G2716->A encodes Valine886->Isoleucine, that has not been previously reported in human C3, although Ile is found in this position in mouse and rat C3.

Our sequence includes start and stop codons, with a complete signal sequence and should, therefore, encode functional C3.

20 Levels of up to 1.7 μ g/ml expressed wild type C3 in culture supernatants of COS-1 cells (transfected using lipofectamine and the pcDNA3 (Invitrogen) expression vector) have been detected by ELISA. No detectable C3 was produced by cells transfected with pcDNA3 vector alone. Furthermore, analysis of the expressed product by cleavage reactions followed by immunoprecipitation, SDS-PAGE and immunoblotting demonstrated that:

30 (i) the primary translation product had been correctly processed into the mature two-chain form;

(ii) this product was, like native C3, cleavable to C3b by C3 convertase (CVFBb); and

(iii) the expressed protein was, like native C3, not cleavable by factor H plus I, but became cleavable after conversion to C3b by C3 convertase enzyme. This confirms that our starting plasmid can be translated into functional C3.

For an alternative description of a construction and expression of a C3 coding sequence see reference [25].

EXAMPLE 1: Production of C3 that has the arginine residues at both factor I cleavage sites (amino acid positions 1303 and 1320) converted to glutamine residues to prevent cleavage of the C3b fragment by factor I.

a) Mutagenesis

Mutagenic oligodeoxynucleotides used were QRI1 (caactgcccagccaaagctccaagatcacc), QRI2 (gccagcctcctgcaatcagaagagaccaag), and AFL4149 (taataaattcgaccttaaggtcaccataaaac), as well as the corresponding antisense oligodeoxynucleotides -QRI1n (ggatgatcttggagctttggctgggcagttg), QRI2n (cttggtctcttctgattgcaggaggctggc) and AFL4149n (gttttatggtgaccttaaggtcgaatttatta).

QRI1 and QRI1n specify the replacement of arginine for glutamine at the factor I cleavage site at amino acid residue 1303 in the C3 precursor sequence (by changing G3968C3969 to AA in the cDNA sequence), and QRI2 and QRI2n effect the same substitution at the factor cleavage site at amino acid residue 1320 (by changing nucleotide G4019 to A).

AFL4149 and AFL4149n introduce a cleavage site for the

restriction endonuclease AflIII at position 4149 in the cDNA sequence (by changing C4149 to T) without altering the encoded amino acid sequence. These two primers were used as markers, allowing successful mutagenesis to be identified on the basis of cleavage of the DNA product by AflIII.

Mutagenesis was effected using the 'gapped plasmid' method. A batch of PGC3 ('UPGC3'), enriched in uridine in place of thymidine, was prepared by growth in *E. Coli* strain CJ236 in the presence of 0.25 µg/ml uridine. This plasmid was digested with *Sma*I and the 7.2kb product ('US1') agarose gel purified to remove a 0.5kb fragment from the C3 sequence (residues 1463-1947). The other component of the gapped plasmid ('DN2') was prepared by digesting PGC3 with *Dra*III plus *Nae*I and purifying the 5.1kb piece twice by agarose gel electrophoresis. 200ng DN2 was mixed with approximately 500ng US1 in 50 µl H₂O, heated to 100°C and cooled slowly to below 50°C, before adding 20 µl to 25 µl of 2XT7 buffer (100mM Tris/HCl/pH 7.4/ 14mM MgCl₂, 100mM NaCl, 2mM dithiothreitol, and 1mM each of ATP, dATP, dCTP, dTTP and dGTP) plus 10nmol of each 5'-phosphorylated mutagenic primer (one reaction used QRI1, QRI2 plus AFL4149, another reaction used QRI1n, QRI2n plus AFL4149n). The mixtures were reheated to 70°C for 5 min and cooled slowly (over 30-60 min) to 20°C. At 0°C, 10 units of T7 DNA polymerase plus 80 units T4 DNA ligase are added. The mixture (total volume 50 µl) was incubated first at 0°C, for 5 min, then at room temperature for 5 min, and finally at 37°C for 3 hours. 1 µl of each mixture was used to transform 100 µl supercompetent XL1 *E. Coli* (Stratagene) according to the manufacturer's instructions.

Ampicillin resistant colonies were screened for *Afl*III cleavage, and successful mutants were grown up in 100ml cultures from which the plasmids were isolated and sequenced (using a sequencing primer C3pa-3876, cttcatggtgttccaagcct, matching nucleotides 3876-3895 of C3 cDNA) to characterise mutations at the factor I cleavage sites.

For an alternative protocol for "gapped plasmid" mutagenesis see references [26,27].

b) Transfer of mutant DNA to eukaryotic expression vector

The C3 coding fragments from mutant plasmids were excised by double digestion with *Hind*III and *Nae*I. *Dra*I was also included to incapacitate the residual plasmid. The C3 coding sequence was agarose gel purified and ligated into pcDNA3 vector (Invitrogen) that had been linearised with *Hind*III and *Eco*RV enzymes and dephosphorylated with calf intestinal phosphorylase. Ligation mixtures were used to transform supercompetent XL1 *E. coli*, which were then plated onto culture plates containing ampicillin.

A random selection (three or four) of ampicillin resistant colonies were grown up in 2-3ml cultures and small scale isolation of the plasmid DNA. The plasmids containing the correct insert were identified by digestion of the plasmid DNA with restriction endonucleases *Eco*RI, *Hind*III and *Afl*III. The corresponding colonies grown up in 100ml cultures and the plasmids purified by the standard procedure. These mutants were originally constructed from PGC3 and so retained the two ATG's 5' to the coding region. This region (plus the 5'

3kb of the C3 coding sequence) was therefore excised with *HindIII* plus *EcoRI* and replaced by ligation of the same segment cut out of PC3. These reconstructed vectors were prepared by the standard procedure and used for transfection of COS cells.

c) Expression of wild-type and mutant C3's

Mutants and wild-type C3 were transiently expressed from plasmids transfected into COS-1 cells using lipofectamine® (GIBCO) according to the manufacturer's instructions. Typically, $1-1.5 \times 10^5$ cells per well of a standard 6 well culture plate were transfected with 2-4 µg of plasmid using 9 µl of lipofectamine reagent. Supernatants were assayed for C3 secretion, and typical yields of 0.3-1.7 µg per ml supernatant were obtained 3-6 days after transfection.

Results

a) Generation of mutants

The following mutants, named according to the mutagenic oligodeoxynucleotide sequences that have been incorporated, have so far been isolated:-

(i) 3 mutants with both QRI1 and QRI2 mutations plus AFL4149: C3M-26, C3M-58 and C3M-61;

(ii) 1 mutant with QRI1 and QRI2 but without AFL4149: C3M-8; and

(iii) 1 mutant with QRI2 and AFL4149, but without QRI1: C3M-51 (used in example 3)

b) Validation that functional effects were due to the mutations specifically introduced at the factor I cleavage sites

5 Sequencing has confirmed the absence of other alterations in 178-350 bases around the mutated region of each mutant. The sequence of one mutant produced by this procedure, C3M-51 (see example 3), has been analysed throughout the entire 'gap' (bases 2463-5067) used in
10 mutagenesis, and no other deviations from the wild-type sequence were found.

Furthermore, representative sequencing of a total of 2922 bases from all mutants have not revealed any single point
15 mutations that could have been caused by polymerase-mediated errors. The expressed mutants all displayed the two-chain structure and cleavage by C3 convertases characteristic of native C3. In summary, the mutants used are unlikely to contain any unwanted changes although
20 they have not been completely re-sequenced.

EXAMPLE 2: Production of C3 that has the arginine residue at one factor I cleavage site (amino acid position 1303) converted to a glutamine residue

25 The procedure of Example 1 was followed except that only mutagenic oligodeoxynucleotides AFL4149 plus QRI1 or AFL4149n plus QRI1n (i.e. no QRI2 or QRI2n), were used in mutagenesis.

30

Results

a) Mutants obtained

2 mutants with QRI1 and AFL4149 but without QRI2 were

isolated:-C3M-I23,27. The mutant C3M-I23 was expressed, as described in Example 1.

5 This protein was cleavable by CVFBb. The C3b-like product was relatively (compared to the wild-type) resistant to cleavage at position 1303 by factors I and H, but could still be cleaved at position 1320. This C3b derivative is therefore partially resistant to factor I.

10 EXAMPLE 3: Production of C3 that has the arginine residue at one factor I cleavage site (amino acid position 1320) converted to a glutamine residue

15 The procedure of Example 1 was followed except that only mutagenic oligodeoxynucleotides AFL4149 plus QRI2 or AFL4149n plus QRI2n (i.e. no QRI1 or QRI1n), were used in mutagenesis. In addition, the method used in example 1 also yielded one mutant with QRI2 and AFL4149, but without QRI1.

20

Results

a) Mutants obtained

25 3 mutants with QRI2 and AFL4149 but without QRI1 were isolated:-C3M-51, C3M-Q2, C3M-Q13. The mutant C3M-51 was expressed, as described in Example 1. This protein was cleavable by CVFBb. The C3b-like product was not readily cleaved at position 1320 by factors I and H, but it could still be cleaved at position 1303. This C3b derivative is
30 therefore partially resistant to factor I.

EXAMPLE 4 Analysis of the functional effects of mutations.

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

NaN₃, and once in 20 mM Tris/HCl, 137 mM NaCl, 0.1% (v/v) Tween 20, pH 7.6 before eluting in 1% SDS/2% 2-mercaptoethanol (90-100°C, 5 min). These eluates were separated by SDS-PAGE, electroblotted onto nitrocellulose and the C3 bands detected by probing with affinity-purified sheep anti-human C3 immunoglobulin followed by horse radish peroxidase-coupled donkey anti-sheep immunoglobulin (Sigma) and detection using the "Enhanced Chemiluminescence" substrates supplied by Amersham. A photograph of a 2 minute exposure to X-ray film is shown. The visible C3-derived bands are indicated by labelled arrows, and the individual samples (1-4, A-D) are those just described. (The prominent band of about 50 kDa (between the 46 and 68 kDa bands) present in all samples is the heavy chain of the IgG used in the immunoprecipitation and detected by the horse radish peroxidase-coupled donkey anti-sheep immunoglobulin.).

Results (see Figure 3)

1. All the untreated samples (1-A, 2-A, 3-A, 4-A) contain bands of the correct migration for alpha and beta chains of C3, indicating that all the mutants are expressed, and post-translationally processed correctly. The presence of 43 or 46 kDa bands in these samples indicates the presence of some factor H + factor I-like activity in the culture medium. Spontaneous hydrolysis of C3 during the 3 day biosynthetic period produces C3i which is cleaved by this activity. In the unmutated C3 this generates bands of 43 kDa and 75 kDa (the 75 kDa band is invisible because (i) it is hidden by the 75 kDa beta chain, and (ii) the antibody used to develop the western blot has very little activity towards this portion of the C3 alpha chain:- its presence was

subsequently confirmed by reprobing with a rat monoclonal antibody, "Clone-3", that is specific for this region). The addition of factors H and I without CVFBb (1-C, 2-C, 3-C, 4-C), did not cleave the remaining C3 indicating
5 that this represented active C3 (thiolester intact).

2. The unmutated C3 (1) is cleaved by CVFBb and the C3b product is further cleaved by endogenous enzymes in 1-B or added factors H and I in 1-D. The 43 kDa band
10 indicates cleavage at Arg¹³²⁰, and the 68 kDa band (visible in longer exposures) indicates cleavage at Arg¹³⁰³

3. The mutant C3M-123 (Arg¹³⁰³->Gln) was cleavable by
15 CVFBb and the product was relatively resistant to endogenous factor H and I-like activity (2-B), with distinct amounts of alpha' chain (C3b) persisting, but was still cleavable when extra factor H and I were added (2-D). The 43 kDa product indicates cleavage at Arg¹³²⁰,
20 (a faint band at 71 kDa representing the other fragment of the alpha' chain could be seen in longer exposures) but no 68 kDa band was present, showing that this mutant is resistant to cleavage at the mutated Gln¹³⁰³.

4. The mutant C3M-26 (Arg¹³⁰³->Gln, Arg¹³²⁰->Gln) was
25 cleavable by CVFBb and the C3b-like product (alpha') was resistant to endogenous factor H and I-like activity (3-B). It was also very resistant to the additional factors H and I (3-D) in comparison with the unmutated C3 (1) and
30 other mutants (2 and 4). There was a small amount of 46 kDa product indicating some cleavage at the mutated Gln¹³⁰³ (the accompanying 68 kDa fragment was also visible on longer exposures). There was little or no detectable 43 kDa that would correspond to any cleavage at Gln¹³²⁰.

Therefore the Arg->Gln mutation at position 1303 is less effective than that at position 1320 at preventing cleavage by factor I. (This slow residual cleavage might also be occurring in the mutant C3M-I23 (Arg¹³⁰³->Gln),
5 but the 46 kDa intermediate is probably being rapidly processed to 43 kDa by further cleavage at the unmutated Arg¹³²⁰.)

5. The mutant C3M-51 (Arg¹³²⁰->Gln) was cleavable by
10 CVFBB and the product was cleaved by endogenous factor H and I-like activity (4-B), and by additional factor H and I (4-D). The 46 kDa product (and faint 68 kDa band) indicates cleavage at Arg¹³⁰³. However, the absence of a
15 43 kDa band indicates that it is not cleaved at the mutated Gln¹³²⁰.

Example 5 Comparison of various amino acid substitutions
at position 1303

20

1. Introduction

The previous examples described mutations of arg 1303 and arg 1320 to glutamine residues. Both mutations imparted
25 resistance to cleavage at those positions by factor I. However, there was a small but detectable degree of cleavage at gln 1303. Therefore a number of other amino acid substitutions at this position have been made and tested. Cleavage occurs, in decreasing order of efficacy
30 when residue 1303 is: Arg > Tyr > [Cys or Trp] > Gln > [Glu or Gly]. These results are unexpected because (i) all known naturally occurring human factor I-mediated cleavages occur C-terminal to arginine residues, so it would have been deduced that the enzyme had a requirement

for arginine; and (ii) if it did cleave at other residues one would predict that they would have to be electrostatically similar to arg, i.e. a basic residue (lys or his), (e.g. trypsin selectively cleaves C-terminal to arg, lys or his), so one could not have predicted cleavage of the tyrosine substitution.

Therefore substitution of arg 1303 with glycine or glutamic acid is preferred for the purpose of creating a derivative of C3 resistant to inactivation by factor I.

2. Methods

2.1 Mutagenesis: the degenerate mutagenic primer used was:

caactgcccagc(gt)(ag)(cg)agctccaagatcacc (letters in brackets indicate mixture of bases at that position).

Mutants were constructed either by the gapped-plasmid method (as described in the earlier examples), or by the "megaprimer method" (V. Picard et al, *Nuc Acid Res* 22:2587-91, (1994)), in which the upstream primer was caccaggaactgaatctagatgtgtccctc and the downstream primer was gttttatggtgaccttaagggtcgaatttatta. All mutations were performed on templates in which the C3-encoding DNA had already been mutated such that amino acid residue 1320 was glutamine, and a restriction site for AflII had been introduced at position 4149 (as described in the earlier examples) and were confirmed by DNA sequencing.

2.2 Expression: mutants were expressed in COS cells using the pcDNA3 vector as described in the earlier examples, biosynthetically labelled with [³⁵S]methionine

in serum-free medium.

2.3 Assay: the supernatants were treated with CVFBB
(formed by reaction of CVF with factors B and D in
5 magnesium-containing buffer) and factors H and I followed
by immunoprecipitation with anti-C3 and separation by
SDS-polyacrylamide gel electrophoresis performed under
reducing conditions (as described in the earlier
examples). The gel was fixed, treated with Amersham
10 "Amplify" reagent, dried and exposed to autoradiography
film to yield the result shown in the figure.

3. Results

15 Factor I-mediated cleavage at position 1303 (site 1),
without cleavage at 1320 (site 2) (where this has been
mutated to glutamine) produces bands of 46 and 68 kDa.
It can be seen that cleavage occurs in the order:
20 arg(R) > tyr(Y) > cys(C) and trp(W) > gln(Q) > gly(G) and
glu(E). The wild-type (arginine at both positions) is
cleaved at both positions to produce fragments of 43 (too
small to be visible on this gel) and 68 kDa.

25

4. Figure

The results are shown in Figure 4. The residues at site 1 (position 1303) and site 2 (1320) are indicated above the respective tracks.

Example 6 Demonstration of enhanced resistance to inactivation by factors I and H after mutation of arg 1303 to gln

1. Introduction

The earlier examples demonstrated that conversion of either arg 1303 or arg 1320 to glutamine made that site resistant to cleavage by factor I. Mutation of both sites makes a molecule that is resistant to cleavage at either site. Here, we further demonstrate that mutation of arg 1303 to gln alone (without alteration to arg 1320) results in a considerable resistance, compared to the wild-type, to functional inactivation by factors I and H.

2. Method

2.1 Expression: The preparation of the arg 1303->gln mutation was described in an earlier example. This was transfected into CHO (a common laboratory cell line derived from chinese hamster ovary cells) by the calcium phosphate method, and stable transfectants selected on the basis of resistance to G418 ("Geneticin" available from Sigma). Cell culture supernatants were collected, and the expressed C3 was partially purified by sodium sulphate precipitation (10-20% (w/v) fraction), and ion-exchange chromatography on Q-sepharose and mono-Q

sepharose (A W Dodds *Methods Enzymol* 223: 46 (1993)).

2.2 Assay: Sheep erythrocytes were coated with SO16 monoclonal antibody (R A Harrison and P J Lachmann *Handbook of Experimental Immunology* 4th Edition chpt. 39 (1986)) and 4.4 ml of a 5% (v/v) suspension was then incubated with approximately 10 μ g C2, 24 μ g C4 and 1 μ g C1 (purified human components) for 10 min at 37°C in CFD (R A Harrison and P J Lachman *supra*). 0.8 ml of this mixture was then incubated for 105 min with 0.25 ml containing the semi-purified mutant or wild-type C3 and EDTA to a final concentration of 12.5 mM. The cells were then washed in CFD and used in CFD containing 0.1% (w/v) gelatin (CFD-gel). Radioligand binding with [125 I]-labelled clone 4 monoclonal anti-C3 antibody was used to confirm that similar amounts of wild-type or mutant C3b were deposited.

For the assay, 40 μ l of a 5% suspension of cells was diluted in 250 μ l CFD-gel and 50 μ l aliquots were incubated with 50 μ l CFD-gel containing dilutions of factors I and H to final concentrations of 100, 10, 1 and 0 μ g/ml each, at 37°C for 30 min. 0.9 ml of CFD was then added, the cells pelleted by centrifugation and washed twice more with 1 ml of CFD each time. The cells were then resuspended in 100 μ l CFD-gel containing 100 μ g/ml factor B, 100 μ g/ml properdin, 1 μ g/ml factor D and 0.3 mM NiCl_2 . After 10 minutes at 37°C, 0.9 ml of CFD containing 10 mM EDTA and 2% (v/v) normal guinea-pig serum. After a further 30 min at 37°C, unlysed cells were pelleted by centrifugation, and the degree of lysis determined by measuring the absorbance of the supernatant at 412 nm. The absorbance equivalent to 100% lysis was determined from an aliquot of cells lysed in water, and

hence the percentage lysis was calculated.

This assay measures the ability of deposited C3b to form a functional C3bBbP convertase. Conversion to iC3b prevents convertase formation and subsequent lysis in serum/EDTA.

3. Results

10

15

20

The result shown in the figure indicates that more than ten times as much factor I and factor H are required to abrogate the hemolytic activity of the arg 1303->gln mutant, when compared to the wild-type. This mutation is therefore advantageous for the creation of a derivative of C3 whose C3b product is resistant to inactivation by factors H and I. The effect could either be due to the greater resistance to cleavage at position 1303 (when arg is mutated to gln), or to greater resistance to cleavage at position 1320 when cleavage can first take place at position 1303.

4. Figure

25

The results are shown in Figure 5. The x-axis indicates the concentration of factors H and I. Q1 represents the arg 1303->gln mutation. % lysis is measured as described in the methods.

Discussion

The essential features of Human C3, with respect to modified variants described herein are as follows:-

5

(i) The molecule has a functionally C3b-like derivative in that it can combine with functionally active human factor B, which can then be cleaved by human factor D to form an enzyme capable of cleaving human C3.

10

(ii) The amino acid sequences of derivatives are more homologous to C3 from humans than to C3 from any other species for which a sequence is presently known, or to any other presently known protein sequence. Structural features of C3 present in wild-type protein, but not necessarily in modified derivatives, include the following:-

15

(a) The DNA coding sequence and translated protein sequence for the variant of human C3 used in the examples of the invention described herein are given in Figures 2 and 1 respectively. This protein sequence differs from the published sequence [2] at just two amino acids (details are given in the examples). It is assumed that many more variations are compatible with C3 function, even though most will not be present in the population.

20

25

(b) The primary translation product is proteolytically processed into two disulphide-linked chains, alpha (residues 672-1663) and beta (residues 23-667), with removal of the signal sequence (residues 1-22).

30

(c) The mature protein contains a thiolester bond between residues Cys1010 and Gln1013.

(d) C3 convertases cleave C3 to remove C3a (residues 672-748). This reaction is followed by breakage of the thiolester bond.

- 5 (e) In the presence of factor H, factor I cleaves C3b between residues Arg1303 and Ser1304, and between Arg1320 and Ser1321.

Modifications made to the native C3 molecule

10

Replacement of Arg1303 by Gln

This modification is at one site of cleavage of C3b by factor I. The effect is to reduce the rate of cleavage by
15 factor I at this position. The change to glutamine was selected to take away the positive charge of the arginine, which is likely to be important for the serine protease activity of factor I, while retaining a hydrophilic character and a similar side-chain size that
20 should minimise any disruptions to the tertiary protein structure. Evidence supporting this presumption is that the mutation did not prevent processing into a two-chain structure, formation of a thiolester or cleavage of C3 by C3 convertase. Mutation of Arg1303 to another amino acid
25 can achieve a similar or even a superior effect, as demonstrated in Example 5.

It may also be possible to reduce this cleavage by mutating Ser1304 (the other side of the cleavage site) or
30 other residues involved in the enzyme-substrate interaction.

Replacement of Arg1320 by Gln

This modification is at the other site of cleavage of C3b by factor I. The effect is to drastically reduce (virtually abolish) the rate of cleavage by factor I at this position. The change to glutamine was made on the same criteria described above, and this mutation also did not prevent processing into a two-chain structure, formation of a thiolester or cleavage of C3 by C3 convertase. Again, mutation to another amino acid may achieve the same effect, as may mutation of Ser1321 or other residues involved in the enzyme-substrate interaction.

When in combination the two mutations, Arg1303-Gln and Arg1320-Gln, protect the C3b from inactivation and hence maintain its ability to form part of an active C3bBb convertase. Other mutations (including combinations of mutations) that abolish both cleavage reactions could also be used (for example Arg 1303 Glu or Arg 1303 Gly could be used in combination with Arg 1320 Gln).

EXAMPLE 7 Various mutations that reduce the interaction
of C3b/C3i with factor H

7.1 Introduction

5 Other laboratories have produced evidence based either on
the effects of synthetic peptides (Ganu, V.S. and Muller-
Eberhard, H.J., 1985, *Complement* 2:27; Becherer, J.D. et
al., 1992, *Biochemistry* 31: 1787-1794), or limited
10 mutagenesis (Taniguchi-Sidle, A. and Isenman, D.E., 1994
J. Immunol. 153: 5285-5302) to suggest that the residues
752-761 in the primary sequence of the C3 transcript (see
figure 1) could be involved in the interaction with
factor H. However, other published evidence suggests
15 that only residues 767-776 are involved in the
interaction with factor H, whereas residues 752-761 are
important for the interaction with factor B (Fishelson,
1991, *Mol. Immunol.* 28:545-552). We surmised that more
extensive mutagenesis of this region might reduce the
20 affinity for factor H and therefore be desirable for the
objective of creating a C3 derivative that is resistant
to factor H. Furthermore, we guessed that the important
residues to mutate could be the prominent acidic residues
(aspartic and glutamic acids) and that it would be
25 desirable to change them to neutral residues less likely
to mediate strong interactions. In this example we
changed residue 752-754 from Asp-Glu-Asp to Gly-Ser-Gly,
in combination with changing residues 758-760 from Glu-
Glu-Asn to Gly-Ser-Gly. The product displayed reduced
30 cleavage characteristics consistent with a reduction in
the susceptibility to factor H. This provides evidence
that C3 can be modified to reduce the binding of factor
H, and hence the susceptibility to factors H and I.
These modifications are desirable for the creation of a

C3 convertase that is stable under physiological conditions.

7.2 Method

5

The methods of mutagenesis, expression and analysis have been described in the earlier examples. The mutagenic oligonucleotide that was synthesised had the sequence:
agtaacctgggttcgggcatcattgcaggatcgggcatcgtttcc.

10

7.3 Results

The results of cleavage reactions are shown in Figure 6. These indicate that:

15

1. Addition of CVFBb to wild-type C3 results in elimination of the alpha chain (track 2) because the C3b that is formed is susceptible to the low concentrations of factor I and H in the culture supernatant. C3i that has been formed during expression or this subsequent incubation has been broken down to iC3i in the same way. Addition of exogenous factors I and H (tracks 3 and 4) are therefore no different from tracks 1 and 2 respectively, because the medium itself contains sufficient factor H and I activity to effect complete cleavage.

25

2. In contrast, treatment of the mutant C3 with CVFBb (track 6) does not result in disappearance of the alpha chain. There is some generation of alpha', corresponding to C3b, but some or all of this remains, indicating that the persistence of alpha chain is not merely the result of a failure of cleavage by CVFBb. The remaining uncleaved alpha chain in track 2 may therefore represent

30

C3i that has not been cleaved by the endogenous activities of factors H and I, although it is also possible that some of this represents native C3 persisting if the mutant has acquired a partial resistance to CVFBb. Addition of high concentrations of exogenous factors H and I (track 7 and 8) does produce depletion of alpha and alpha' chains, indicating that (i) the mutant is not completely resistant to these factors, and (ii) the alpha chain uncleaved by CVFBb in track 2 is predominantly derived from C3i (which is cleavable by factors H and I but not by CVFBb) rather than from native C3 (which is cleavable by CVFBb but not by factors H and I). Still not all the alpha chain is cleaved, even in track 8, probably because of the resistance to factors H and I.

Therefore mutation of residues 752-754, and residues 758-760 can generate a C3 molecule that can still be cleaved by C3 convertases, but is partially resistant to the actions of factors H and I. In view of other published data, this is most probably because the mutations have modified a region that is involved in the interaction with factor H and hence have resulted in a reduced affinity for factor H.

EXAMPLE 8 A site in C3 that can be mutated to modify the interaction of C3i with factor B

8.1 Introduction

The previous examples have demonstrated that mutations to C3 can modulate the interactions with factors H and I. In order to discover other sites in C3 that might

interact with factor B, we compared the known sequences of C3 molecules from different species, as well as with available sequences for C4 and other homologous proteins. We identified the region corresponding to residues 1427-1433 of human C3 that might be involved in C3 and C4 specific functions. This could include interaction with factor B (or its homologue, C2, in the case of C4), but not necessarily because other potential functions include thiolester formation, conversion into C3b (or C4b form), interaction with substrate C3 and/or C5 in convertase activity and interaction with factor I and its cofactors. Therefore selected residues were mutated to the corresponding residues (based on sequence alignments) found in another homologous protein, in this case human C5. Thus residue 1427 was changed from an Arg to a Gln, residue 1431 from a Lys to Asp, and residue 1433 from a Glu to a Gln. The resulting mutant was found to be susceptible to cleavage by C3 convertase (CVFBb) and the C3b product was cleavable by factors H and I. However, this mutant did not support the conversion of factor B to Bb plus Ba, which is dependent on the binding of factor B to C3i (or C3b). Therefore we have evidence that mutation of this region has diminished the interaction with factor B. Whilst this is undesirable for the generation of a super-active C3 convertase, it does provide an indication that other modifications to this region of C3 will also alter the interaction with factor B, and some of these will probably increase the affinity. As a consequence such mutations may also increase the stability and activity of the bimolecular convertase enzyme, C3bBb (or C3iBb).

8.2 Methods

The alignments shown in Table 1 overleaf illustrate why we considered that this region was a candidate for mutagenesis. We surmised that characters of certain residues were well conserved in C3 and C4 but distinctly different in the other proteins. Residues 1427, 1431 and 1433 were selected because their charged nature might be indicative of groups involved in protein-protein interactions. The changes were made to the corresponding residues in human C5 because these displayed very different electrostatic properties, but within the context of some other conserved residues that might indicate a similar local structure.

Table 1
Alignments of sequences of C3 and related molecules for region of residues 1427-1435 of human C3

Protein	Species	RESIDUE (Human)									
		1427	1428	1429	1430	1431	1432	1433	1434	1435	
C3	Human	R	Y	I	S	K	Y	E	L	D	
	Mouse	R	Y	I	S	K	Y	E	M	N	
	Rat	R	Y	I	S	K	Y	E	M	D	
	G. pig	R	Y	I	S	K	Y	E	L	D	
C4	Rabbit	R	Y	I	S	K	Y	E	L	N	
	Cobra	R	Y	I	S	K	F	E	I	D	
	Xenopus	K	Y	I	S	K	Y	E	V	N	
	Trout	R	Y	I	E	K	F	E	M	D	
slp	Human	R	Y	V	S	H	F	E	T	E	
	Mouse	R	Y	V	S	H	F	E	T	D	
	Mouse	R	Y	V	S	H	F	E	T	D	
C3/C4-like	Hagfish	N	Y	I	V	Q	Y	E	I	R	
	Lamprey	K	Y	I	S	N	Y	E	I	T	
C5	Human	Q	L	F	T	D	Y	Q	I	K	
	Mouse	Q	L	L	T	D	Y	Q	I	K	
A2M	Human	P	T	V	K	M	L	E	R	S	
	Mouse	P	S	V	K	R	L	Q	D	Q	
	Rat	P	T	V	K	M	L	E	R	S	
PZP	Human	P	T	V	K	M	L	E	R	S	
Murinoglobulin	Mouse	P	T	V	K	K	L	E	R	L	
	Rat	P	S	V	K	K	L	Q	D	Q	
A1M	G. Hamster	P	T	V	K	K	L	E	R	S	
A1I3	Rat	P	T	V	K	K	L	E	R	L	

The methods of mutagenesis, expression and analysis of C3 cleavage reactions were as described in the earlier examples (Examples 1-4). The mutagenic oligonucleotide was synthesised with the sequence:

5 tgggtgttgaccaatacatctccgactatcagctggacaa.

Assay for turnover of factor B.

The expressed product was purified from the COS cell medium by affinity purification on a column of Clone-3-Sepharose as described in Example 9. This method results in considerable conversion of the thiolester broken form, C3i. Wild-type C3 was isolated by the same procedure. Dilutions of the wild-type C3 (1/5, 1/25 and 1/125) were run on an SDS-PAGE gel (reducing conditions) along with the mutant C3, and silver staining indicated that the mutant was present at a concentration equivalent to slightly less than the 1/25 but much more than the 1/125 dilution of wild-type. The same dilutions were used in the assay of factor B turnover. 5µl of these C3's were incubated with 25µl of CFD-G containing 5µg/ml factor D and approximately 1.6µg/ml of ¹²⁵I-labelled factor B (approx. 1000-2000 dpm/µl) for 3h at 37°C. The samples were then analysed by SDS-PAGE (reducing conditions) with autoradiography of the dried gel. The results are shown in Fig. 7.

8.3 Results

As shown in Fig. 7, distinct cleavage of factor B occurs even at a 1/125 dilution of the wild-type C3 (C3i). In contrast, no significant cleavage was observed in the presence of the mutant C3, even undiluted which should be at a concentration higher than the 1/125 sample of the wild-type.

This mutant therefore appears to have an impaired ability to support the cleavage of factor B, most likely due to a reduction in its binding affinity for factor B. Therefore this is a region of C3 that can be mutated to modulate the interaction between C3i (or C3b) and factor B and perhaps also the stability of the convertase (C3iBb or C3bBb).

10 EXAMPLE 9 Purification of expressed mutant C3 molecules

9.1 Introduction

This example demonstrates how the mutant C3 molecules may be isolated from an expression medium, such as the culture medium of transfected eukaryotic cells. By simple affinity purification the C3 molecules are obtained in sufficient purity for functional tests and for conjugation to antibody by the method described in Example 10. Although elution from an antibody is accompanied by hydrolysis of a considerable proportion of the internal thiolester, the C3i product is still a suitable precursor for the generation of an active C3 convertase, as well as for the production of C3i-antibody conjugates. This approach is also likely to be useful as part of the preparation required for *in vivo* use.

9.2 Method

30 Affinity-purification on Clone-3-Sepharose.

Clone-3 is a rat monoclonal antibody that is specific for C3 and its derivatives, including C3b and C3i (Lachmann, P.J. et al., 1980, *J. Immunol.* 41:503-515). Other monoclonal antibodies against C3 are available, and in

some cases have been successfully used to isolate C3 from small quantities of human plasma (Dodds, A.W., 1993, *Methods Enzymol.* 223:46-61) and are therefore also likely to be applicable for the isolation of molecules expressed ex vivo. The IgG fraction was coupled to Sepharose CL-4B using cyanogen bromide (methodology may be found in Harrison and Lachmann, 1986, *Handbook of Experimental Immunology*, 4th edn., Ed.s Weir, Herzenberg, Blackwell and Herzenberg; Blackwell, Oxford). Culture supernatants were either passed directly through a column of this resin (re-circulated), or first concentrated by precipitation with 25% (w/v) Na_2SO_4 , and resolubilization and dialysis into PBS, 5 mM NaN_3 . The column is then washed successively with (i) PBS, 5 mM NaN_3 and (ii) PBS containing 1 M NaCl. Bound C3 elutes with 50 mM Na borate buffer, pH 10.5, and is immediately neutralised by collection of 0.9 ml fractions into 0.1 ml 1 M Tris/HCl pH 7. The material is then dialysed into PBS, 5 mM NaN_3 .

Preparation of C3 bearing a "His-Tag"

A "His-Tag" is a string of histidine residues that displays affinity for columns bearing Nickel ions. This method has been employed to aid the isolation of expressed proteins. We thought that this could be useful for the isolation of expressed mutant C3 molecules so we have used insertion mutagenesis to generate a plasmid encoding C3 with a tail of 6 histidine residues at the carboxy terminus (immediately carboxy-terminal to residue 1663). This location for the his tag was selected so as to minimise interference with the synthesis, folding, processing and disulphide bond formation of the nascent C3. Residue 1661 is a cysteine residue that is involved in a disulphide bond to a residue earlier in the sequence (probably Cys 1537; Dolmer, K. and Sottrup-Jensen, L.,

1993, *FEBS-Lett* 315: 85-90) and therefore it seemed prudent to make the insertion beyond this structural feature. The mutation was introduced using the "gapped-plasmid" technique used in Example 1, using the mutagenic oligonucleotide synthesised with the sequence:

5 tgggtgcccccaaccatcatcatcatcattgaccacaccccc.

Incorporation of the correct sequence was confirmed by DNA sequencing. This DNA sequence may now be transferred to an expression vector. After transfection of eukaryotic cells, it should be possible to isolate the expressed C3 by affinity for a column bearing Nickel ions, or by any other matrix with specific affinity for the "His-Tag".

15 9.3 Results

A number of mutant C3 have been purified on the Clone-3-Sepharose, including those described in Examples 1 and 2 expressed in CHO cells. The products retained the ability to support the cleavage of factor B by factor D. The same method was used to isolate the mutant described in Example B2, expressed in COS cells. Silver-staining of SDS-PAGE gels indicated that the isolated products were not 100% pure, but often appeared to be greater than or equal to 50% pure. This comes from starting materials generally containing less than 10µg/ml C3 in 10% (v/v) fetal calf serum plus other cellular proteins. In addition the C3's were not degraded during isolation, and endogenous factor H and I activity appeared to have been removed.

Purification by virtue of the "His-Tag" involves milder elution conditions from a column bearing Nickel ions.

For example, EDTA has been used. Application of this method to C3 should therefore allow isolation without rupture of the internal thiolester bond.

5

EXAMPLE 10 Conjugation of C3i to antibody and use to target C3 convertase activity against a particular cell

10 *10.1 Introduction*

One aspect of the invention is that stable C3 convertases derived from mutant C3 molecules will cause enhanced C3 conversion which, if localised at a particular target site, will promote complement-dependent attack of that target. The favoured approach for targeting the response is to couple the mutant C3 molecule, as either the C3i or C3b derivative, to an antibody specific for the desired target. In this example we demonstrate a working methodology for formation of such conjugates, which is applicable to mutant C3i or C3b molecules and can be used on material affinity-purified from an expression system, even if the thiolester of C3 has been broken in the process. By coupling C3i to an antibody that specifically binds to sheep erythrocytes, we further show that the the conjugate fixes C3i to the erythrocyte surface such that a convertase, C3iBbP, can be formed that initiates lysis of these cells when other complement components are supplied in the form of normal guinea-pig serum (in EDTA to prevent de-novo formation of C3 convertases). Hence conjugation to antibody can be used to target a C3i molecule to initiate complement-dependent attack of a particular cell type. This example uses wild-type C3i, from human plasma, that forma a C3

convertase in vitro. In vivo, wild-type C3i and C3b are broken down by factor H and I. Therefore a mutant C3, constructed according to the plans in this patent to be resistant to factors H and I and therefore forming a stable C3 convertase, would be advantageous in a physiological context.

10.2 Method

10 (i) Generation and purification of C3i-antibody conjugate

The antibody used was the IgG fraction isolated from a polyclonal rabbit anti-sheep erythrocyte antiserum. 1.1 mg was incubated with 75 nmol of SPDP in conjugation buffer, pH 7.5 (20 mM KH_2PO_4 , 60 mM Na_2HPO_4 , 0.12 M NaCl) for 2h at room temperature. The PDP-IgG was purified by gel-filtration on a Superose-6 column (Pharmacia) (in a phosphate buffer, pH 7.4, containing 0.5 M NaCl). Reduction of a sample with dithiothreitol was used to estimate 4 PDP groups coupled per molecule of IgG. C3i was prepared by treatment of purified C3 with 0.1 M methylamine, pH 7.2 (2h at 37°C). Excess methylamine was removed by gel-filtration followed by dialysis into conjugation buffer. 18 nmole of C3i was mixed with 1.7 nmoles of PDP-IgG in 1.26 ml conjugation buffer and incubated for 1 day at room temperature followed by 1.5 days at 4°C. Figure 8 shows a Coomassie Blue stained SDS-PAGE gel of the conjugation reaction mixture showing the appearance of a species of approximately 350 kDa that was not present in either PDP-IgG or C3i. This species was partially purified by gel-filtration on the Superose-6 column in a phosphate buffer, pH 7.4, containing 0.5 M NaCl and then dialysed into PBS. It eluted before the C3, in a volume from which a molecular weight of 300-400

kDa could be estimated by calibration with globular molecular weight standards. Concentrations of conjugate, free antibody and uncoupled C3 were estimated from a Coomassie-stained SDS-PAGE gel (non-reducing conditions).
5 Two-dimensional SDS-PAGE (first dimension unreduced, second dimension reduced) revealed a pattern compatible with a 1:1 conjugate between IgG and C3i.

(ii) Demonstration that the C3-antibody conjugate can be
10 used to target convertase activity against a particular cell.

20µl of dilutions of the conjugate (0 (no conjugate), 1/100, 1/50, 1/10) were incubated with 100µl of approximately 1% (v/v) sheep erythrocytes (prewashed in
15 CFD) for 1 hour at 37°C. Parallel incubations were performed with equivalent amounts of PDP-IgG (no C3) and C3 alone. The cells were then washed 4 times in CFD and resuspended to 100µl in CFD-G. 50µl of this were lysed with 150 µl H₂O, followed by addition of 800 µl of CFD
20 containing 10 mM EDTA and 2% (v/v) NGPS. The other 50µl of conjugate-coated cells were incubated for 15 min at 37°C with 50µl of CFD-G containing 190µg/ml factor B, 2µg/ml factor D, 20µg/ml properdin and 0.6 mM NiCl₂, followed by lysis with 900µl of CFD containing 10mM EDTA
25 and 2% (v/v) NGPS. After 30 min at 37°C, the cells were pelleted by centrifugation (2000 X g, about 3 min) and the optical absorbance of the supernatant was measured at 412 nm. Using the H₂O-treated samples as 100% lysis, and a buffer blank devoid of cells, the % lysis was
30 calculated, as shown in Fig. 9. The conjugate produced dose-dependent lysis, whereas neither the PDP-IgG nor the C3i alone generated any lysis significantly above that observed in the absence of any such treatment.

10.3 Summary of Results

The method used has proved successful for coupling C3i to IgG as shown by:

5

1. The formation of a band of appropriate size (about 350 kDa) for a 1:1 C3:IgG conjugate shown by SDS-PAGE in Fig. 8.

10

2. Two-dimensional SDS-PAGE (first dimension non-reduced, second dimension reduced) indicated that this species contained both IgG and C3i.

15

3. The elution characteristic of this species on gel-filtration is again consistent with a molecule of about 350 kDa.

20

4. The conjugate displays a haemolytic activity that is not displayed by either PDP-IgG or C3i (Fig. 9).

The haemolytic assay (Fig. 9) further demonstrates that:

25

1. The specific anti-sheep erythrocyte antibody has localised the C3i to the target cell (sheep erythrocyte) membrane, preventing it from being removed by washing (in contrast to free C3i).

30

2. The conjugate retains the activity of the C3i in that it is still able to form a C3 convertase by reaction with properdin and factors B and D.

3. This convertase can initiate complement-dependent attack of the target, in this case by activating the lytic pathway (C5-9) to lyse the erythrocyte.

Additional data from other laboratories show that cobra venom factor can be coupled to an antibody and that these conjugates can target complement activation against a particular cell type (Vogel, 1988, *Targeted. Diagn. Ther.*, 1:191-224; Muller, B. and Muller-Ruchholtz, W., 1987, *Leuk. Res.* 11:461-468; Parker, C.J., White, V.F. and Falk, R.J., 1986, *Complement* 3:223-235; Petrella, E.C. et al, 1987, *J. Immunol. Methods* 104:159-172). These data support the contention that C3 modified so that it is capable of forming a stable C3 convertase, like cobra venom factor, could be used to target complement-mediated responses, as outlined in this invention.

EXAMPLE 11 Demonstration that mutant C3 molecules induce factor B turnover in normal human serum

11.1 Introduction

A major purpose of the invention described herein is the consumptive depletion of complement activity from biological fluids. The invention describes methods for the manufacture of C3 molecules that are resistant to down-regulation by factors H and I. In this state they will bind factor B and generate active C3 convertases. The activity of these convertases is demonstrated by the haemolytic assay employed in Example 6. Such a convertase will therefore consume C3. If the convertase is unstable, it will dissociate without much C3 conversion. However this will allow binding of fresh factor B, and its conversion to Bb and Ba. Thus the mutant C3 will promote the consumption of factor B,

leading ultimately to the disablement of the alternative pathway, and its inability to amplify classical pathway stimulation. If a stable C3 convertase is formed, turnover of factor B will be reduced, but consumption of C3 will be increased. Both situations can therefore be desirable. In this example we demonstrate that mutant C3 molecules that are modified to make them resistant to factor I, but without any modification to modify the stability of the convertase, promote accelerated turnover of factor B in human serum. Wild-type C3, in contrast, causes no significant turn-over, presumably because wild-type C3i is rapidly degraded by factors H and I.

11.2 Method

The Mutants prepared are as follows:

Q1R2 Arg1303 changed to Gln (Example 2)

Q1Q2 Arg1303 changed to Gln, plus Arg1320 changed to Gln (Example 1)

E1Q2 Arg1303 changed to Glu, plus Arg1320 changed to Gln (Example 5)

These mutants were all expressed in CHO cells and then purified by precipitation with Na_2SO_4 , followed by affinity purification on Clone-3-Sepharose, as described in Example B3. Wild-type C3 (R1R2) was similarly isolated. By SDS-PAGE with silver-staining, the concentration of Q1 was between 1/5 and 1/25 of the wild-type, the concentration of Q1Q2 was about that of 1/5 wild-type, and the concentration of E1Q2 was between 1/25 and 1/125 of wild-type. All preparations probably contained a majority of thiolester-broken molecules (C3i).

10µl of these C3 preparations were incubated with 10µl of a solution of 20% (v/v) normal human serum in PBS containing 1 mM MgCl₂ and approximately 300 ng ¹²⁵I-labelled factor B (approx. 2-300,000 dpm) for 1 hour at 37°C. 5µl was then analysed by SDS-PAGE (reducing conditions). The dried gel was exposed to autoradiography film to indicate the positions of the bands corresponding to the intact factor B and its cleavage products. These were then excised and counted to accurately determine the degree of cleavage. The value obtained in buffer alone was subtracted as background (encompassing not only background cleavage, but also degradation products and other impurities present in the radioligand preparation).

15

11.3 Results

The resulting degrees of factor B cleavage are shown below:

20

1/25 Wild-type	1.49%
1/5 Wild-type	2.74%
Q1R2	6.19%
Q1Q2	7.41%
E1Q2	6.42%

25

Therefore the factor I resistant mutants all produce greater levels of factor B cleavage than equivalent amounts of wild-type C3 (C3i). With larger doses or longer incubations, complete incapacitation of the alternative pathway should result.

30

The abbreviations used in the foregoing examples include: CFD, complement fixation diluent (defined in Harrison and

- 61 -

Lachmann, 1986, *Handbook of Experimental Immunology*, 4th edn., Ed.s Weir, Herzenberg, Blackwell and Herzenberg; Blackwell, Oxford); CFD-G, CFD containing 0.1% (w/v) gelatin; PBS, phosphate-buffered saline; NGPS, normal guinea-pig serum; SDS-PAGE, SDS-polyacrylamide gel electrophoresis; SPDP, N-Succinimidyl-3-[2-pyridyldithio]propionate.

EXAMPLE 12 Mutation of residues 992-1000

1. **Introduction**

Other laboratories have produced evidence based on the effects of synthetic peptides (Ganu, V.S. and Muller-Eberhard, H.J., 1985, *Complement* 2:27; Becherer, J.D. et al., 1992, *Biochemistry* 31:1787-1794; Fishelson, Z., 1991, *Molecular Immunology* 28:545-552; Lambris, J.D., Ganu, V.S., Hirani, S. and Muller-Eberhard, H.J., 1988, *J. Biol. Chem.* 263:12147-12150) to suggest various residues in human C3b that might be involved in the interaction with Factor H. We have used the different approach of sequence comparison to predict residues involved in C3-specific functions. Site-directed mutagenesis has been performed and has indicated that most of these candidates have little or no influence on the functional susceptibility to Factor H. However, a few mutations did reduce the susceptibility to Factor H. These mutations were made to parts of the molecule that have not previously been identified as interacting with Factor H or modulating its binding. Hence mutagenesis of these defined residues can be used to produce mutant derivatives of C3 that are partially or completely resistant to inhibition by Factor H within a physiological environment, and will form complex C3 convertase enzymes (C3bBb etc) that are similarly resistant to inactivation by Factor H.

Factor H is structurally homologous to other complement inhibitory proteins, including CR1, MCP and DAF. In view of this apparent evolutionary relationship, and mutual competition for binding, it is likely that they interact with C3b in a structurally similar manner to Factor H (Farries et

- 62/1 -

al., 1990, Complement Inflamm. 7:30-41). Therefore the mutations described which modulate the susceptibility to Factor H are also likely to be useful for modulating the interactions with these other proteins, especially for the purpose of evading their complement down-regulatory activities. They may also find application for the modification of the interaction with the SCR domains in Factor B (and the homologous domains in C2 involved in binding to C4b). Mutations to the corresponding regions of C4 and C5 might also be useful to modify their interactions with the SCR domains in C1r and C1s (C4), C4b-binding protein (C4b), and C6 and C7 (C5b).

2. Method

2.1 Searching for residues involved in C3-specific functions.

These predictions were made from alignments of human C3 with all the homologous proteins for which sequences were available through public data bases. These included the functionally equivalent molecules in mouse, rat, guinea-pig, rabbit, cobra, xenopus, chicken and trout, human and mouse C4, human and mouse C5, C3-like proteins from lamprey and hagfish, cobra venom factor (CVF), and human alpha-2-macroglobulin and its homologues. Searches were then made for residues that were conserved among different C3s, but distinctly different in homologues (notably C5 and CVF) that lack the C3-specific functions of interest. Some of these have been mutated to encode the corresponding residues in C5 or CVF, expressed in COS cells and the secreted products tested for cleavage in the presence of CVFBb and Factor H and I. All methods are as described in the standard methods and example 1. A summary of the results is shown in Table II.

- 62/2 -

2.2 Construction and analysis of mutant DV-1AM

This mutant was made in the same way as the other mutants, using the "megaprimer method" as described by V. Picard et al., 1994, *Nuc. Acid. Res.* 22:2587-2591. The mutagenic primer had the sequence ccagatgacaagtgctgccgtcagccagtcagggctgaagcacc encoding the mutations E992S, D993A, D996S, A997Q, E998S and R999G. The up-stream primer had the sequence tgtcatcgtagccgctaaaga (corresponding to deoxynucleotides 2754-2773), and the down-stream primer had the sequence gttttatggtagaccttaagggtcgaatttatta (complementary to deoxynucleotides 4130-4165, with the introduction of a cleavage site for the restriction enzyme Afl II at position 4149). The mutated DNA fragment was ligated into a vector that contained the coding sequence for C3 also with the introduced site for Afl II at position 4149, by cutting both pieces with Afl II and EcoRI (cuts at position 2997), purifying the desired products and ligating together using T4 DNA ligase. Plasmid DNA was isolated from transformed bacterial colonies, and genuine mutants identified by DNA sequencing. At this point it was found that the DNA sequence had been additionally mutated to encode the mutation L1000M. The resulting expression vector was transfected into COS cells, and the secreted expressed product analysed for cleavage reactions as previously described.

3. Properties of mutant DV-1AM

Analysis of the expressed product with the DV-1AM mutations is shown in Figure 10. The points to note are:

- (i) The western blot is developed with monoclonal antibodies to the C3dg region of C3 that detect the precursor, alpha, alpha', 77 and 68 kDa fragments, but not the beta, 43 or 46 kDa fragments.

- 63/1 -

(ii) All bands of the DV-1AM product (lanes B1-4) appear slightly below the equivalent wild-type bands (lanes A1-4). The shift in mobility is a consequence of the mutations made.

(iii) Cleavage of wild-type C3 with CVFBb produces some alpha' chain from C3b, but a larger amount of 68 kDa fragment resulting from cleavage of the C3b to iC3b by endogenous Factor H and I activity (lane A3). Addition of exogenous H and I completes the conversion of C3b to iC3b (lane A4). In contrast, cleavage of the DV-1AM product by CVFBb produces a larger amount of alpha' chain and only a small amount of 68 kDa fragment (lane B3). Addition of exogenous H and I then converts this C3b into iC3b (lane B4). Therefore the mutant C3b is much more resistant than the wild-type to endogenous H and I, although resistance is not complete as indicated by susceptibility to cleavage by high concentrations of exogenously added H and I.

4. Conclusion

The DV-1AM mutation creates resistance to Factor H-dependent cleavage by Factor I. The mechanism is not certain, although because the modified residues are far (in the primary structure) from the sites of cleavage by Factor I, it is likely that it is the interaction with Factor H that is impaired. In this case the mutation will also impart resistance to the other inhibitory activities of Factor H, namely:- (i) competition with Factor B for binding to C3b (or C3i), and (ii) accelerated dissociation of the C3bBb and C3bBbP convertases. The DV-1AM mutation has modified residues either directly or indirectly involved in maintaining the affinity for Factor H. Many different mutations of the

- 63/2 -

same residues will presumably have similar effects, and mutation of only some of the residues modified here, as well as other residues within this segment of the primary structure, is also likely to achieve such an effect.

EXAMPLE 13 Mutation of residues 1001-1005

1. Introduction

As described in the above example, the residues 1001, 1002 and 1005 were also identified as candidates that might be essential for C3-specific functions. Mutation confirms that modification of these residues can be used to impart resistance to Factor H.

2. Method

2.1 Construction and analysis of mutant DV-1B

The method used was as described for the preceding example, with the exception that the mutagenic primer had the sequence aacggctgaacatattaattcataccccctcgggc encoding the mutations K1001N, H1002I and V1005H. Sequence analysis of isolated plasmid DNA confirmed that the correct mutation had been introduced. No other mutations were detected.

3. Properties of mutant DV-1B

Analysis of the expressed product with the DV-1B mutations is shown in Figure 11. The points to note are:

- 64/1 -

(i) The western blot is developed with a polyclonal antibody to the C3 that detects the precursor, alpha, alpha', beta, 43 and 46 kDa fragments strongly, and only weakly detects the 77 and 68 kDa fragments. Note that the alpha and alpha' chains are not transferred and detected with 100% efficiency, so the intensity of these bands is less than expected and a poor guide to the actual amounts present.

(ii) Cleavage of wild-type C3 with CVFBb produces a small amount of alpha' chain from C3b, but most of this is lost due to cleavage of the C3b to iC3b by endogenous Factor H and I activity (lane B3). This appears mostly as 43 kDa fragment, although a small amount of the 46 kDa intermediate is visible. Addition of 2 µg/ml exogenous H (lane B4), with Factor I, causes marked further cleavage and 10-50 µg/ml exogenous H (lanes B5, B6) completes the conversion of C3b to fully cleaved (no alpha' or 46 kDa bands) iC3b. Cleavage of the DV-1B product by CVFBb also produces a small amount of alpha' chain (lane A3; the total amount of C3 present is much less, and the alpha and alpha' bands are very faint). Significantly, the amount of 43 kDa chain generated in the absence of exogenous H and I is less than in the wild-type, and the appearance of the 46 kDa intermediate fragment is relatively greater, indicating less effective cleavage. Addition of exogenous H and I (lanes A4-6) completes the conversion of this C3b into iC3b, but the 43 kDa product is seen to increase dose-dependently upto 50 µg/ml H (lane A6), when the 46 kDa intermediate is still evident. Therefore the mutant C3b is more resistant than the wild-type to endogenous and exogenous H and I, although resistance is not complete as indicated by susceptibility to cleavage by high concentrations of

exogenously added H and I.

4. Conclusion

The DV-1B mutation creates resistance to Factor H-dependent cleavage by Factor I. The mechanism is not certain, although because the modified residues are far (in the primary structure) from the sites of cleavage by Factor I, and the effect was dependent on the dose of Factor H added, it is likely that it is the interaction with Factor H that is impaired. In this case the mutation will also impart resistance to the other inhibitory activities of Factor H, namely:- (i) competition with Factor B for binding to C3b (or C3i), and (ii) accelerated dissociation of the C3bBb and C3bBBP convertases. The DV-1B mutation has modified residues either directly or indirectly involved in maintaining the affinity for Factor H. Many different mutations of the same residues will presumably have similar effects, and mutation of only some of the residues modified here, as well as other residues within this segment of the primary structure, is also likely to achieve such an effect.

EXAMPLE 14 Mutation of residues 1152-1155

1. Introduction

As described in the above examples, the residues 1152, 1153 and 1155 were also identified as candidates that might be essential for C3-specific functions. Mutation confirms that modification of these residues can be used to impart resistance to Factor H.

2. Method

- 65/1 -

2.1 Construction and analysis of mutant DV-6

The method used was as described for the preceding example, with the exception that the mutagenic primer had the sequence atctcgctgcgcaaggctttcgatatttgcgag encoding the mutations Q1152R, E1153K and K1155F. Sequence analysis of isolated plasmid DNA confirmed that the correct mutation had been introduced. No other mutations were detected.

3. Properties of mutant DV-6

Analysis of the expressed product with the DV-6 mutations is shown in Figure 11. The points to note are:

(i) As described in the preceding example, the western blot is developed with a polyclonal antibody to the C3 that detects the precursor, alpha, alpha', beta, 43 and 46 kDa fragments strongly, and only weakly detects the 77 and 68 kDa fragments. Note that the alpha and alpha' chains are not transferred and detected with 100% efficiency, so the intensity of these bands is less than expected and a poor guide to the actual amounts present.

(ii) Cleavage of wild-type C3 with CVFBb produces a small amount of alpha' chain from C3b, but most of this is lost due to cleavage of the C3b to iC3b by endogenous Factor H and I activity (lane B3). This appears mostly as 43 kDa fragment, although a small amount of the 46 kDa intermediate is visible. Addition of 2 µg/ml exogenous H (lane B4), with Factor I, causes marked further cleavage and 10-50 µg/ml exogenous H (lanes B5, B6) completes the conversion of C3b to fully cleaved (no alpha' or 46 kDa bands) iC3b. Cleavage of the DV-6 product by CVFBb also produces a small amount of alpha' chain (lane C3). Significantly, the amount of 43 kDa

- 65/2 -

chain generated in the absence of exogenous H and I is less than in the wild-type, and the amount of 46 kDa intermediate is relatively greater, indicating less effective cleavage. Addition of exogenous H and I (lane C4-6) completes the conversion of this C3b into iC3b. However, whereas with the wild-type the 46 kDa intermediate was eliminated by 10 µg/ml H (lane B5), indicating complete cleavage, this species still persisted with the mutant with H at this concentration (lane C5), and complete cleavage was only apparent when 50 µg/ml H was used. Therefore the mutant C3b is more resistant than the wild-type to endogenous H and I, although resistance is not complete as indicated by susceptibility to cleavage by the highest concentrations of exogenously added H and I.

4. Conclusion

The DV-6 mutation creates resistance to Factor H-dependent cleavage by Factor I. The mechanism is not certain, although because the modified residues are far (in the primary structure) from the sites of cleavage by Factor I, and the effect was dependent on the dose of Factor H added, it is likely that it is the interaction with Factor H that is impaired. In this case the mutation will also impart resistance to the other inhibitory activities of Factor H, namely:- (i) competition with Factor B for binding to C3b (or C3i), and (ii) accelerated dissociation of the C3bBb and C3bBbP convertases. The DV-6 mutation has modified residues either directly or indirectly involved in maintaining the affinity for Factor H. Many different mutations of the same residues will presumably have similar effects, and mutation of only some of the residues modified here, as well as other residues within this segment of the primary structure, is also likely to achieve such an effect.

TABLE 1: SUMMARY OF EFFECTS OF MUTATIONS ON SUSCEPTIBILITY TO FACTOR H

Mutation	Amino Acid Changes	Inhibition of Factor H dependent cleavage by Factor I
CV-2	E776K	-
CV-1	P963K, P964A, A965R, D966K	-
DV-1AM	E992S, D993A, D996S, A997Q, E998S, R999G, L1000M	++
DV-1B	K1001N, H1002I, V1005H	+
DV-3	T1031G, E1032N, Q1033H, E1035N, K1036I	-
DV-4	V1070K, K1071G, R1072G, A1073S, P1074A	-
CV-5	R1134Q	-
DV-6	Q1152R, E1153K, K1155F	+
DV-7N	D1174N	-
DV-9	D1216G, K1217E, N1218D, R1219H	-
CV-4	R1260N, G1264E	-
RY-1	R1427Q, K1431D, E1433Q	-

Key: -, no inhibition detected; +, small inhibition; ++, larger inhibitory effect

- 67/1 -

EXAMPLE 15 Alteration of residues 1546-1663**1. Introduction**

Unlike previous examples (12-14) modification of residues 1546-1663 was not based on consideration of sequence comparisons between C3 and related proteins. Instead the modification described was created by accident, a consequence of an unintended nucleotide deletion that caused a frame-shift in the translation of the C-terminal residues. The resulting product displayed considerable resistance to Factor H-dependent cleavage by Factor I. Therefore similar modifications created by design are likely to be useful for conferring resistance to the regulatory actions of Factor H and/or Factor I.

2. Method

A vector equivalent to NC3, but carrying additional mutations to 3151g, 3152g, 3154a, 3156c, 3159c, 3163a, 3165t, 3167t, 3168t that translate into the amino acid changes T1031G, E1032N, Q1033H, E1035N and K1036I was digested with restriction enzymes Pvu I (cuts in vector sequence) and BsrG I (cuts at nucleotide 4692), and the 6.1 kb band isolated by agarose gel electrophoresis. Another vector equivalent to NC3 but carrying the insertion of catcatcatcatcatcat after nucleotide 5049, to encode the insertion of amino acids HHHHHH at the C-terminus, was similarly digested with Pvu I and BsrG I, and the 4.4 kb fragment isolated. These two DNA fragments were ligated together, and a complete plasmid was isolated. DNA sequencing found that a single nucleotide (a4696 or a4697) had been lost. The predicted consequence is that amino acids 1546-1663 cannot be

- 67/2 -

translated in frame. Instead, there will be 48 residues read out of the normal frame until a stop codon is reached.

The product, "HDV-3X", was expressed in COS cells and tested as described in the preceding examples.

3. Properties of mutant HDV-3X

Analysis of the expressed product with the HDV-3X modification is shown in figure 12. The points to note are:

(i) The western blot is developed with monoclonal antibodies to the C3dg region of C3 that detect the precursor, alpha, alpha', 77 and 68 kDa fragments, but not the beta, 43 or 46 kDa fragments.

(ii) HDV-3X is compared with mutant DV-3, as the equivalent product without the additional C-terminal modification. The HDV-3X displays a smaller alpha chain, but normal sized 68 kDa and 77 kDa products, consistent with a truncation at the C-terminus.

(iii) Cleavage of DV-3 C3 with CVFBb in the presence of Factor I produces some alpha' chain from C3b, but a larger amount of 68 kDa fragment resulting from cleavage of the C3b to iC3b dependent on endogenous Factor H (lane A2). Addition of exogenous H completes the conversion of C3b to iC3b (lane A3-5). The conversion is virtually complete with only 1 µg/ml Factor H (lane A3). In contrast, cleavage of the HDV-3X product by CVFBb produces a larger amount of alpha' chain and

- 68/1 -

only a small amount of 68 kDa fragment (lane B2). Addition of exogenous H is ineffective in converting this C3b into iC3b (lane B3-5). Only slight formation of 68 kDa and 77 kDa products is detectable even with 25 µg/ml H (lane A5). Therefore the HDV-3X C3b is much more resistant than the wild-type to endogenous H and I. The fact that resistance is partially overcome by higher amounts of Factor H suggests that the affinity for Factor H may be greatly reduced.

4. Conclusion

The HDV-3X modification creates resistance to Factor H-dependent cleavage by Factor I. The mechanism is not certain, although because the modified residues are far (in the primary structure) from the sites of cleavage by Factor I, it is likely that it is the interaction with Factor H that is impaired. In this case the mutation will probably also impart resistance to the other inhibitory activities of Factor H, namely: - (i) competition with Factor B for binding to C3b (or C3i), and (ii) accelerated dissociation of the C3bBb and C3bBbP convertases. The HDV-3X modification has affected residues either directly or indirectly involved in maintaining the affinity for Factor H. Many different deletions or mutations of the same residues will presumably have similar effects, and deletion or mutation of only some of the residues modified here is also likely to achieve such an effect.

The HDV-3X modification was not created by design. But the methods of choice for creating this and related modifications are likely to be specific methods of site-directed mutagenesis, including those methods described in preceding examples.

- 68/2 -

EXAMPLE 16 Modification of residues 954 and 955 to prevent Factor I mediated cleavage at this site.

1. Introduction

Previous mutagenesis at the P₁ residues (1303 and 1320) provided resistance to cleavage by Factor I at the first two sites (examples 1-6). It was not known if prevention of cleavage at these two sites would also prevent cleavage at a third site responsible for release of C3c from C3dg. This third cleavage, which is normally dependent on CR1 (a membrane bound receptor that has been engineered into a soluble form, sCR1) as a cofactor, is relatively slow and has only previously been observed on iC3b (or iC3i) (i.e. after cleavage at sites 1 and 2) and not on C3i or C3b. To test this, the E1Q2 mutant (described in example 11), which is highly resistant to cleavage at sites 1 and 2, was used. If this mutant was still susceptible to cleavage at site 3, it would indicate that it would be desirable to mutate this site to prevent degradation of the molecule in physiological fluids. However, there are conflicting reports in the literature as to whether the cleavage occurs exclusively at the 954-955 bond (Davis, A.E. 3d. Harrison, R.A. & Lachmann, P.J., 1984, *J. Immunol.*, 132:1960-6), or whether cleavage can also occur at other positions, such as 959-960 (Harrison, R.A. et al., 1996 *Molecular Immunology* 33, Suppl. 1, 59, abstract 235; Ekdahl, K.N., Nilsson, U.R. & Nilsson, B., 1990, *J. Immunol.* 144: 4269-74). Initially we mutated residue 954 from arginine to Glutamic acid (to make E1Q2E3) because (i) this appears from the above publications to be the P₁ residue of one of the cleavage sites, and (ii) from example 5 at site 1,

- 68/3 -

where mutation to Glutamic acid imparted higher resistance to cleavage than other substitutions. In addition other mammalian species (mouse, rat, guinea pig, rabbit) of C3 have Glutamine and Glycine at the residues equivalent to 954 and 955, instead of the arginine and Glutamic acid of human C3 (e.g. Mavroidis, M., Sunyer, J.O. & Lambris, J.D., 1995, *J. Immunol.* 154:2164-2174). These data suggest that this site (954-955) would not be well cleaved in other species, and that another site, such as 959-960, might be more important (Harrison, R.A., et al., 1996, *Molecular Immunology* 33, Suppl. 1, 59, abstract 235). The equivalent mutations of arg954 to Gln, and Glu955 to Gly were therefore made to human C3 to make the E1Q2QG3 mutant.

2. Method

The method used for mutant construction was as described for preceding examples, with the exception that the mutagenic primer for the E1Q2E3 mutant had the sequence `gaacgcctggggcgaagaaggagtgcag` encoding the mutation R954E, and the mutagenic primer for the E1Q2QG3 mutant had the sequence `aacgcctggggccaaggaggagtgcagaa` encoding the mutations R954Q, E955G. The product was ligated into a construct that contained the mutations encoding E1Q2 (E1303, Q1320, as described in examples 5 and 11). Sequence analysis of isolated plasmid DNA confirmed that the correct mutation had been introduced. No other mutations were detected. The resulting expression vectors were transfected into COS cells, and the secreted expressed product analysed for cleavage reactions as previously described.

- 68/4 -

3. Factor I-mediated cleavages of E1Q2, E1Q2E3 and E1Q2QG3 mutants

Analysis of the expressed products is shown in figure 13. The points to note are:-

- (i) The western blot is developed with monoclonal antibodies to the C3dg region of C3 that detect the precursor, alpha, alpha', 77 and 68 kDa fragments, but not the beta, 43 or 46 kDa fragments. In addition the 86 kDa product of cleavage at site 3, without cleavage at sites 1 or 2, will be detected.
- (ii) The figure shows that the 86 kDa product is indeed formed by Factor I-mediated cleavage of E1Q2 in the presence of sCR1 (lane A3), but not when Factor H is the cofactor (lane A2).
- (iii) The 86 kDa product is not formed in either of the E1Q2E3 (C) or E1Q2QG3 (B) mutants, even in the presence of sCR1 (C3 and B3).

4. Conclusion

(i) Factor I-mediated cleavage at site 3 can still occur when cleavage at sites 1 and 2 have been blocked. Therefore additional blockage of cleavage at site 3 is desirable to prevent degradation of any mutant product that is otherwise only resistant at sites 1 and 2, when used in a physiological environment.

(ii) Cleavage at site 3 can be blocked by mutation of residue 954 to Glu, and by mutation of 954 and 955 to Gln and Gly. Therefore other mutations of residues 954 and/or 955 are also likely to impart resistance to cleavage at site 3.

- 68/5 -

(iii) The mutations shown did not allow cleavage at other putative positions of third site cleavage (such as 959-960), even though such sites were not mutated. This would indicate that either 954-955 is the only significant site of cleavage, or that other cleavages require prior cleavage at 954-955, or that mutation of these residues prevents cleavage at other positions by a different mechanism (such as conformational distortion). In any case, mutations of 954 and/or 955 are effective means of preventing degradation of C3b or C3i-like products.

- 68/6 -

EXAMPLE 17 Modifications to the carboxy-terminal region of C3 that inhibit cleavage by Factor I.

1. Introduction

Example 15 provided evidence that mutation or deletion of residues 1546-1663 imparted resistance to cleavage by Factor I. This example provides further smaller scale mutants that also impart this resistance, as well as various mutations that do not.

2. Method

The method used for mutant construction was as described for preceding examples, with the exception that the mutagenic primers had the sequences shown in table III, encoding the mutations indicated. Sequence analysis of isolated plasmid DNA confirmed that the correct mutation had been introduced. No other mutations were detected. The resulting expression vectors were transfected into COS cells, and the secreted expressed product analysed for cleavage reactions as previously described.

3. Resistance of various mutants to cleavage by Factor I

The results of cleavage assays performed with Factors I and H on these mutants are shown in figures 14 and 15. Note that the western blots are developed with monoclonal antibodies to the C3dg region of C3 that detect the precursor, alpha, alpha', 77 and 68 kDa fragments, but not the beta, 43 or 46 kDa fragments.

- 68/7 -

Figure 14 shows that the wild-type (NC3,A), FT-3(D), FT-4(E) and FT-5(F) products are cleaved by Factor I, as indicated by the appearance of 77 kDa and/or 68 kDa bands and the disappearance of the alpha chain. In contrast FT-1 and FT-2 are not cleaved.

Figure 15 shows that the wild-type (NC3,A), FR-1(B), FR-2(C), FR-3(D) and FR-4(E) products are cleaved, while again the FT-2 product (F) is cleaved.

4. Conclusions

(i) Even the small truncation of FT-2 is sufficient to impart resistance to cleavage by Factor I. Such resistance can therefore be achieved by deletion of some or all of residues 1636-1663. This conclusion is supported by the resistance displayed by FT-1 which includes deletion of residues 1636-1663, with additional deletion/modification of residues 1591-1635.

(ii) As residues 1636-1663 are required for Factor I-mediated cleavage, many other modifications of these residues are likely to generate resistance.

(iii) Not all modifications of these residues impart resistance. Ineffective modifications include those defined by FT-5, FR-1, FR-2, FR-3 and FR-4, as well as the modifications defined by FT-3 and FT-4 that modify residues other than 1636-1663.

(iv) These data imply that the residues within 1636-1663 that are required for cleavage are those that have not been modified by FT-5, FR-1, FR-2, FR-3 or FR-4. Therefore some of the residues 1649-1660 may be critical.

TABLE III. MUTANTS USED IN EXAMPLE 17				
Mutant	Sequence of mutagenic primer	Sequence replaced	Residues	Replaced by
FT-1		REA	1591-1593	TN stop
FT-2		E	1636	stop
FT-3		LSSDFWGE	1607-1614	KEALQI
FT-4		IIGKD	1621-1625	RYIYPLDSL
FT-5		C	1661	S
FR-1		EEDE	1633-1636	RDTT
FR-2		QDEENQKQ	1638-1645	SS
FR-3		QDEENQKQ	1638-1645	RSTRQRAA
FR-4		D	1648	AFLAN

REFERENCES:

1. Bergmann, M. & Fruton, J.S. (1941) *Adv. Enzymol.*, 1:63-98.
- 5 2. de Bruijn, M.H. & Fey, G.H. (1985), *Proc. Natl. Acad. Sci. U.S.A.* 82:708-712
3. Crawford-MH et al. (1988) *Circulation*. 78:1449-58
4. Daha, M.R. & van Es, L.A. (1982) *Immunol.* 43:33-38.
5. Farries, TC; Lachmann, PJ & Harrison, RA (1988)
- 10 *Biochem. J.* 252:47-54
6. Farries, TC; Lachmann, PJ & Harrison, RA (1988)
- Biochem. J.* 253:667-75
7. Forty, J; Hasan, R; Cary, N; White, DJ & Wallwork, J
- (1992) *Transplant. Proc.* 24:488-9
- 15 8. Fritzinger, D.C. et al. (1992) *J. Immunol.*
- 149:3554-3562
9. Harrison, R.A. & Lachmann, P.J. (1980) *Mol. Immunol.*
- 17:9-20.
10. Kalli, K.R., Hsu, P. & Fearon, D.T. (1994) *Springer*
- 20 *Semin. Immunopathol.* 15:417-431.
11. Kinoshita, T; Takata, Y; Kozono, H; Takeda, J; Hong, KS
- & Inoue, K (1988) *J. Immunol.* 141:3895-901
12. McNearney, TA; Odell, C; Holers, VM; Spear, PG;
- Atkinson, JP (1987) *J. Exp. Med.* 166:1525-35
- 25 13. Nicol, P.A.E. & Lachmann, P.J. (1973) *Immunol.*
- 24:259-275
14. Pangburn, MK & Muller-Eberhard, HJ (1984) *Springer*
- Semin. Immunopathol.* 7:163-92
15. Rother, K. & Till, G.O. (eds) (1988) "The complement
- 30 *System"* (Springer-Verlag Berlin Heidelberg, Germany)
16. Van den Berg, C.W., Aerts, P.C. & Van Dijk, H. (1991)
- J. Immunol. Methods* 136:287-294.
17. Vogel, CW; Smith, CA & Muller-Eberhard, HJ (1984) *J.*
- Immunol.* 133:3235-41

18. Weisman, HF et al. (1990) *Science* 249:146-51.
19. Wu, R. (ed.) (1993) *Methods Enzymol.* 217: ch.s 12-14
(Academic Press, San Diego, U.S.A.)
20. Botto, M, Fong, K.Y., So, A.K., Koch, C. & Walport, M.J.
5 (1990) *J. Exp. Med.* 172:1011-7
21. Sambrook, J., Fritsch, E.F. & Maniatis, T. (1989)
"Molecular Cloning. A Laboratory Manual" second
edition (Cold Spring Harbor Laboratory Press)
22. Fishelson, Z. (1991) *Mol. Immunol.* 28:545-52.
- 10 23. Taniguchi-Sidle, A & Isenman, D.E. (1993) *Mol.*
Immunol. 30:54.
24. Lambris, J.D., Avila, D., Becherer, J.D. &
Muller, Eberhard, H.J. (1988) *J. Biol. Chem.*
263:12147-50.
- 15 25. Taniguchi-Sidle, A. and Isenman, D.E. (1992) *J.*
Biol. Chem. 267:635-643.
26. Hofer, B. and Kuhlein, B. (1993) *Methods Enzymol.*
217:173-189.
27. Morinaga, Y., Franceschini, T., Inouye, S. and
20 Inouye, M. (1984) *Bio-technology* 2:636-639.
28. Harrison, R.A. and Lachmann, P.J. (1986) "*Handbook*
of Experimental Immunology" (eds Weir, Herzenberg,
Blackwell and Herzenberg; Blackwell, Oxford) 4th ed.,
29. Kotwal, G., J., and Moss, B., *Nature* (1988) 335
25 (6186):176-8.

CLAIMS:

1. A native complement pathway protein modified such that the protein is capable of forming a down-regulation resistant C3 convertase.
2. A protein as claimed in claim 1 which is a modified human protein.
3. A protein as claimed in claim 1 or claim 2 which is more resistant to cleavage by factor I than the native protein.
4. A protein as claimed in any one of claims 1 to 3 which is a modified C3 protein.
5. A protein as claimed in claim 4 wherein the protein is modified by replacement of either Arg-1303, Arg-1320, or both by another amino acid.
6. A protein as claimed in claim 5 wherein Arg-1303, Arg-1320 or both are replaced by glutamine, tyrosine, cystine, tryptophan, glutamic acid or glycine.
7. A protein as claimed in claim 6, wherein Arg-1320 is replaced by glutamine.
8. A protein as claimed in claim 6 or 7, wherein Arg-1303 is replaced by glutamic acid, glycine or glutamine.
9. A protein according to any preceding claim which has reduced susceptibility to Factor H and/or Factor I relative to native human C3, said protein having one or more amino acid changes relative to native human C3 in

- 72 -

the region corresponding to amino residues 752-754 and/or residues 758-780 of native human C3.

10. A protein according to claim 9 wherein the one or more amino acid changes are changes from acidic amino acid residues to neutral amino acid residues.

11. A protein according to claim 9 or claim 10 wherein the amino acid residue changes are changes from Asp-Glu-Asp to Gly-Ser-Gly.

12. A protein according to any preceding claim having amino acid changes relative to native human C3 at amino acid residues corresponding to residues 1427, 1431 and/or 1433 of native human C3.

13. A protein according to any preceding claim which includes one or more mutations relative to amino acid residues 992-1005 (EDAVDAERLKHLIV) of human C3, such that the C3b and C3i products, or their derived C3 convertases, are resistant to the complement inhibitory activity of Factor H.

14. A protein according to claim 13 which includes one or more mutations relative to amino acid residues 992 (E), 993 (D), 996 (D), 997 (A), 998 (E), 999 (R), 1000 (L), 1001 (K), 1002 (H), 1005 (V) of human C3.

15. A protein according to claim 14 which includes one or more of the following mutations E992S, D993A, D996S, A997Q, E998S, R999G, L1000M, K1001N, H1002I and V1005H.

16. A protein according to any preceding claim which includes one or more mutations relative to amino acid

- 73 -

residues 1152-1155 (QEAK) of human C3, such that the C3b and C3i products, or their derived C3 convertases, are resistant to the complement inhibitory activity of Factor H.

17. A protein according to claim 16 which includes one or more mutations relative to amino acid residues 1152 (Q), 1153 (E) and 1155 (K) of C3.

18. A protein according to claim 17 which includes one or more of the following mutations Q1152R, E1153K and K1155F.

19. A protein according to any of claims 13 to 18 which is resistant to the complement inhibitory activity of CR1, MCP and/or DAF.

20. A protein according to any preceding claim which has one or more amino acid deletions, substitutions or insertions relative to amino acids 1546-1663 of native human C3; wherein said protein has reduced susceptibility to Factor H and/or Factor I, relative to native human C3.

21. A protein according to claim 20 comprising one or more amino acid deletions relative to amino acids 1546-1663 of native human C3.

22. A protein according to claim 20 or 21 comprising a deletion of all amino acids corresponding to amino acids 1546-1663 of native human C3.

23. A protein according to claim 20 comprising one or more different amino acids relative to native human C3 at a region corresponding to amino acid residues 1546-1663

of native human C3.

24. A protein according to claim 23 wherein the amino acids at the region corresponding to amino acids 1546-1663 of native human C3 can result from a frame-shift mutation in DNA encoding said native human C3.

25. A protein according to any of claims 20 to 24 which has one or more amino acid deletions, substitutions or insertions relative to amino acids 1636-1663 of native human C3; wherein said protein has reduced susceptibility to Factor H and/or Factor I, relative to human C3.

26. A protein according to claim 25 comprising one or more amino acid deletions relative to amino acids 1636-1663 of native human C3.

27. A protein according to claim 25 or 26 comprising a deletion of all amino acids corresponding to amino acids 1636-1663 of native C3.

28. A protein according to claim 25 comprising one or more different amino acids relative to native human C3 at a region corresponding to amino acid residues 1636-1663 of native human C3.

29. A protein according to claim 25 comprising one or more amino acid deletions substitutions or insertions relative to amino acids 1649-1660 of native human C3; wherein said protein has reduced susceptibility to Factor H and/or Factor I, relative to human C3.

30. A protein according to any preceding claim which has one or more amino acid deletions substitutions or insertions relative to amino acid residues 954 and/or 955

- 75 -

of native human C3 and which has a reduced susceptibility to cofactor-dependent (e.g. CR1 or Factor H) Factor I-mediated cleavage at this position.

31. A protein according to claim 30 in which the amino acid residue at the position corresponding to residue 954 of human C3 is different from residue 954 of human C3, which protein has reduced susceptibility to cofactor dependent (e.g. CR1 or Factor H) Factor I-mediated cleavage at said position.

32. A protein according to claim 31 in which said amino acid residue is glutamic acid, and which has reduced susceptibility to cofactor dependent (e.g. CR1 or Factor H) Factor I-mediated cleavage at said position.

33. A protein according to claim 30 in which the amino acid residues at positions corresponding to residue 954 and residue 955 of human C3 are glutamine and glycine respectively which protein has reduced susceptibility to cofactor dependent (e.g. CR1 or Factor H) Factor I-mediated cleavage at said position.

34. A protein according to claim 30 in which the amino acid residue at a position corresponding to residue 955 of human C3 is different from residue 955 of human C3, which protein has reduced susceptibility to cofactor dependent (e.g. CR1 or Factor H) Factor I-mediated cleavage at said position.

35. A fragment or a variant of a protein according to any preceding claim, said fragment or variant having C3 convertase activity and also having resistance to the complement inhibitory activity of Factor H, Factor I, CR1, MCP and/or DAF.

- 76 -

36. A DNA sequence coding for a protein as claimed in any one of claims 1 to 34 or for a fragment or a variant as claimed in claim 35.
37. A DNA construct (e.g. a vector) comprising a DNA sequence as defined in claim 36.
38. A protein as defined in any one of claims 1 to 34 or a fragment or variant as claimed in claim 35 for use in therapy.
39. A conjugate comprising a protein as defined in any one of claims 1 to 34 or a fragment or variant as claimed in claim 35 linked to a specific binding moiety.
40. A conjugate as claimed in claim 39 wherein the specific binding moiety is a specific binding protein.
41. A conjugate as claimed in claim 40 wherein the specific binding protein is an antibody or antigen binding fragment thereof.
42. The use of a protein as defined in any one of claims 1 to 34, or a fragment or variant as defined in claim 35, or a conjugate as defined in any one of claims 39 to 41 in the manufacture of a medicament for use in depleting levels of complement pathway protein.
43. The use as claimed in claim 42 wherein the medicament is for use in preventing rejection of foreign matter.
44. The use as claimed in claim 42 wherein the medicament is for use in localising and/or amplifying endogenous complement protein conversion and deposition

at a specific site.

45. A pharmaceutical formulation comprising one or more proteins as defined in any one of claims 1 to 34, or a fragment or variant as defined in claim 35, or a conjugate as defined in any one of claims 39 to 41 together with one or more pharmaceutically acceptable carriers or excipients.

46. A pharmaceutical formulation as claimed in claim 45 which is for use in depleting levels of complement pathway protein.

47. A pharmaceutical formulation as claimed in claim 45 which is for use in preventing rejection of foreign matter.

48. A pharmaceutical formulation as claimed in claim 45 which is for use in localising and/or amplifying complement protein conversion and deposition at a specific site.

49. A method of reducing complement pathway protein in a mammal which comprises administering to the mammal a protein as defined in any one of claims 1 to 34, or a fragment or variant as defined in claim 35, or a conjugate as defined in any of claims 39 to 41.

50. A method as claimed in claim 49 wherein administration occurs using a pharmaceutical formulation as defined in claim 45.

1/19

FIG. 1(I)

10	20	30	40	50	60
MGPTSGPSLL	LLLLTHLPLA	LGSPMYSIIT	PNILRIESEE	TMVLEAHDAQ	GDVPVTVTVH
70	80	90	100	110	120
DFPGKKLVLS	SEKTVLTFAT	NHMGNVTFIT	PANREFKSEK	GRNKFTVQCA	TFGTQVVEKV
130	140	150	160	170	180
VLVSLQSGYL	FIQTDKTIYT	PGSTVLYRIF	TVNHKLLFVG	RTVMVNIENP	EGIPVKQDSL
190	200	210	220	230	240
SSQNGLGVLV	LSWDIPELVN	MGQWKIRAYY	ENSPOQVFST	EFEVKEYVLP	SFEVIVEPTE
250	260	270	280	290	300
KFYTYVNEKG	LEVTITARFL	YGKKVEGTAF	VIFGIQDGEQ	RISLFESLKR	IPFEDGSGEV
310	320	330	340	350	360
VLSRKVLLEG	VQNPRAEDLV	GKSLVVSATV	ILHSGSDMVC	AERSGIPVTV	SPYQIHFTKT
370	380	390	400	410	420
PKYFKPGMPF	DLNVFVTNPD	GSPAYRVFVA	VOGEDTVQSL	TQGGGVAKLS	INTHPSOKPL
430	440	450	460	470	480
SITVTRTKQCE	LSEAEQATRT	MQALPYSTVG	NSNNYLHLSV	LRTELRPGET	LNVNELLRMD
490	500	510	520	530	540
RAHEAKIRYY	TYLIMNKGRL	LKAGROVREP	GQDLVVLPLS	ITDFIPSFH	LVAYYTLIGA
550	560	570	580	590	600
SGQREVVAQS	VWVDVKDSGV	GSLVVKSGQS	EDROPVPGQC	MTLKIEGDHG	ARVVLVAVDK
610	620	630	640	650	660
GTVFLNKKNK	LTOSKIWDVV	EKADIGCTFG	SGKDYAGVFS	CAGLTFTSSS	GQQTAAQRAEL
670	680	690	700	710	720
QCQPPAARRR	RSVOLTEKRM	DKVGKYPKEL	AKCCEDGMRE	NPMRFSCQRA	TPFISLGEAC
730	740	750	760	770	780
KKVFLDCCNY	ITELRROHAR	ASHLGLARSH	LDEDIAEEN	IVSRSEFPES	WLWNVEDLKE
790	800	810	820	830	840
PPKNGISTKL	MNIFLKDSIT	TWEILAVSMS	DKKGICVADP	FEVTVMQDFF	IDLRLPYSVV
850	860	870	880	890	900
RNEQVEIRAV	LYNYRONQEL	KVRVELLHNP	AFCSLATTKR	RHQQTITIPP	KSSLSVPYVI
910	920	930	940	950	960
VPLKTLQCEV	EVKAAVYHHF	ISDGVRKSLK	VVFESIRMNK	TVAVRTLOPE	RLGREGVOKE
970	980	990	1000	1010	1020
DEPPADLSGC	WPDTESETRI	LLOGTFVACM	TEDAVDAERL	MKLIVTFSGC	GEQNMIGMTP
1030	1040	1050	1060	1070	1080
TVIAYVYLDE	TEQWEKFSLE	KROGALELIK	KGYTQCLAFR	QFSSAFAAFV	KRAPSTWLTA
1090	1100	1110	1120	1130	1140
YVVFETSLAV	NLIAIDSOVL	CGAVKWLILE	KOKPDGVTEB	DAPVTRQEMI	GGLRNNNEKD
1150	1160	1170	1180	1190	1200
MALTAFLVLS	LOEAKDICEE	OVNSLFSSIT	KAGDFLEAHY	MNLORSYTVA	IAGYALAQMG

SUBSTITUTE SHEET (RULE 26)

2 / 19

1210	1220	1230	1240	1250	1260
RLKGFLLNKF	LTTAKDKNRW	EDPGKQLYNV	EATSYALLAL	LQLKDFDFVP	PVVRWLNEQR
1270	1280	1290	1300	1310	1320
YYGGGYGSTQ	ATFMVFOALA	QYQKDAPDHO	ELNLDVSLQL	PSRSSKITHR	IHWESASLLR
1330	1340	1350	1360	1370	1380
SEETKENEGB	TVTAEGKGGG	TLSVVVTHYA	KAKDQLTCNK	FDLKVTIKPA	PETETRPQDA
1390	1400	1410	1420	1430	1440
KNTMILEICT	RYRGDQDATH	SILDISMMTG	FAPDTDDLKC	LANGVDRIYS	KYELDKAFSD
1450	1460	1470	1480	1490	1500
RNTLIITYLDK	VSHSEDDCLA	FKVHQYFNVE	LIQPGAVKVY	AYYNLEESCT	RFYHPEKEDG
1510	1520	1530	1540	1550	1560
KLKLCRDEL	CRCAEENCPI	OKSDDKVTL	ERLDKACEPG	VDYVYKTRLV	KVQLSNDFDE
1570	1580	1590	1600	1610	1620
YIMAIECTIK	SGSDEVQVGO	QRTFISPIKC	REALKLEKK	HYLMWGLSSD	FWGEXPNLSY
1630	1640	1650	1660		
IIGKDTWVEH	WPEEDECQDE	ENQKQCQCLG	AFTESMVVFG	CPN	

FIG. 1(II)

3 / 19

FIG. 2(I)

	ccctccct	ctgtccctct	gtccctctga	ccctgcactg	tcccagcacc	
	12	20	30	40	50	60
atgggaccca	cctcaggtcc	cagcctgttg	ctcctgttac	taaccacacc	ccccctggct	
70	80	90	100	110	120	
ctggggagtc	ccatgtactc	tatcatcacc	cccaacatct	tgcggctgga	gagcgaggag	
130	140	150	160	170	180	
accatgggtc	tggaggccca	cgacgcgcaa	ggggatgttc	cagtccactgt	tactgtccac	
190	200	210	220	230	240	
gacttcccag	gcaaaaaact	agtgtctgtcc	agtgagaaga	ctgtgtctgac	ccctgccacc	
250	260	270	280	290	300	
aaccacatgg	gcaacgtcac	cttcacgatc	ccagccaaca	gggagtcca	gtcagaaaaag	
310	320	330	340	350	360	
ggggccaaca	agttcgtgac	cgtgcaggcc	accttcggga	cccaagtgtt	ggagaagggtg	
370	380	390	400	410	420	
gtgtctggca	gcttcagag	cgggtacctc	ttcatccaga	cagacaagac	catctacacc	
430	440	450	460	470	480	
ccctggctcca	cagttctctta	tgggtctctc	accgtcaacc	acaagtctgt	acccgtgggt	
490	500	510	520	530	540	
cggacggcca	tggccaacat	tgagaaccgg	gaaggcatcc	cggccaagca	ggactccttg	
550	560	570	580	590	600	
tcttctcaga	accagctctg	cgtcttgcgc	ttgtcttggg	acattccgga	actcgtcaac	
610	620	630	640	650	660	
atgggtccagt	ggaagatccg	agcctactat	gaaaactcac	cacagcaggt	cttctccact	
670	680	690	700	710	720	
gaatttgagg	tgaaggagta	cgtgtctgccc	agtttcgagg	tcatagtggg	gcctacagag	
730	740	750	760	770	780	
aaattctact	acatctataa	cgagaagggc	ctggaggcca	ccatcacccg	caggttcttc	
790	800	810	820	830	840	
tacgggaaga	aagtggaggg	aactgccttt	gtcatctctg	ggatccagga	tggcgaacag	
850	860	870	880	890	900	
aggatttccc	tgcctgaatc	cctcaagcgc	attccgattg	aggatggctc	gggggaggtt	
910	920	930	940	950	960	
gtctctgagc	ggaaggtaact	gtctgacggg	gtgcagaacc	cccgagcaca	agacctgggtg	
970	980	990	1000	1010	1020	
gggaagtctt	tgtacgtgtc	tgcacccgtc	atcttccact	caggcagtga	catggtgcag	
1030	1040	1050	1060	1070	1080	
gcagagcgca	gggggatccc	catcgtgacc	tctccctacc	agatccacct	caccaagaca	
1090	1100	1110	1120	1130	1140	
cccaagtact	tcaaacccag	aatgcctctt	gacctcatgg	tgttcgtgac	gaaccttgat	
1150	1160	1170	1180	1190	1200	
ggctccccag	ccatccgagt	ccccctggca	gtccagggtc	aggacactgt	gcagtctcta	
1210	1220	1230	1240	1250	1260	

SUBSTITUTE SHEET (RULE 26)

FIG. 2(II) 4/19

acccaggag	atggcggtggc	caaaactcagc	atcaacacac	acccagcca	gaagcccttg
1270	1280	1290	1300	1310	1320
agcatcacgg	tgcgcacgaa	gaagcaggag	ctctcggagg	cagagcaggg	taccaggacc
1330	1340	1350	1360	1370	1380
atgcaggctc	tgccttacag	caccgtgggc	aactccaaca	attacctgca	tctctcagtg
1390	1400	1410	1420	1430	1440
ctacgtacag	agctcagacc	gggggagacc	ctcaacgtca	acttcctcct	gcgaatggac
1450	1460	1470	1480	1490	1500
cgcgcaccag	aggccaagat	cggctactac	acctacctga	tcatgaacaa	gggcaggctg
1510	1520	1530	1540	1550	1560
ctgaaggcgg	gaccccaggt	ggagagagcc	ggccaggacc	tggtagtgct	gccccgtcc
1570	1580	1590	1600	1610	1620
atcaccaccg	acttcacccc	tcccttcggc	ctggtagcgt	actacacgct	gacgggtgcc
1630	1640	1650	1660	1670	1680
agcggccaga	gggaggtggc	ggccgactcc	ctgtgggtgg	acgtcaagga	ctcctggctg
1690	1700	1710	1720	1730	1740
ggctcgctgg	tggtaaaaag	cggccagtca	gaagaccggc	agcctgtacc	tgggcagcag
1750	1760	1770	1780	1790	1800
atgacctga	agatagaggg	tgaccacggg	ggccgggtgg	tactggtagc	cgtggacaag
1810	1820	1830	1840	1850	1860
ggcgtgttcg	tgctgaataa	gaagaacaaa	ctgacgcaga	gtaagatctg	ggacgtggcg
1870	1880	1890	1900	1910	1920
gagaaggcag	acatcggtcg	caccccgggc	agtgggaagg	attacggcgg	tgtcttctcc
1930	1940	1950	1960	1970	1980
gacgcagggc	tgaccttcac	gagcagcagt	ggccagcaga	ccgccagag	ggcagaacct
1990	2000	2010	2020	2030	2040
cagtgcctcc	agccagccgc	ccggcgacgc	cggtccgtgc	agctcacgga	gaagcgaatg
2050	2060	2070	2080	2090	2100
gacaaagtcc	gcaagtaccc	caaggagctc	cgcaagtgcg	ggcaggacgg	catgcgggag
2110	2120	2130	2140	2150	2160
aaccccatga	ggttctcgtg	ccagcgccgg	acccgtttca	tctccctggg	cgaggcgtgc
2170	2180	2190	2200	2210	2220
aagaaggtct	tcctggactc	ctgcaactac	atcacagagg	tcggcgggca	gcacgcgcgg
2230	2240	2250	2260	2270	2280
ggcagccacc	tgggcccggc	caggagtaac	ctggatgagg	acatcattgc	agaagagaac
2290	2300	2310	2320	2330	2340
atcggtttcc	gaagtgaagt	cccagagacc	tggctgtgga	acgttcgagga	cttgaaagag
2350	2360	2370	2380	2390	2400
ccaccgaaaa	atggaatctc	tacgaagctc	atgaatatat	ttttgaaaga	ctccatcacc
2410	2420	2430	2440	2450	2460
acgtgggaga	ttctggctct	gagcatgtcg	gacaagaaaq	ggatctgtgt	ggcagacccc
2470	2480	2490	2500	2510	2520
ctcgaggtca	cagtaatcca	ggactttctc	atccacctgc	ggctacccca	ctctgttctt
2530	2540	2550	2560	2570	2580

FIG. 2(III) 5/19

cgaaacgagc	aggtggaaat	ccgagccgtt	ctctacaatt	accggcagaa	ccaagagctc
2590	2600	2610	2620	2630	2640
aaggcgaggg	tggaactact	ccacaatcga	gccttctgca	gcctggccac	caccaagagg
2650	2660	2670	2680	2690	2700
cgccaccagc	agaccataac	catccccccc	aaqtcctcgt	tgctccgttc	atatgtcacc
2710	2720	2730	2740	2750	2760
gtcccgctaa	agaccggcct	gcaggaagtg	gaactcaagg	ctgctgtcta	ccatcatttc
2770	2780	2790	2800	2810	2820
atcagtgacg	gtgtcaggaa	gtccctgaag	gtcgtgcccg	aaggaatcag	aatgaacaaa
2830	2840	2850	2860	2870	2880
acctgtggctg	ctccacacct	ggatccagaa	cgcttgggcc	gtgaaggagt	gcagaaagag
2890	2900	2910	2920	2930	2940
gacatccccc	ctccagacct	cagtacccaa	gtcccgagca	ccgactctga	gaccagaaat
2950	2960	2970	2980	2990	3000
cccccgcaag	ggacccagct	ggcccgagatg	acagaggatg	cggtcgaccc	ggaacgggctg
3010	3020	3030	3040	3050	3060
aaqcacccca	ctgtgacccc	ctcgggctgc	ggggaacaga	acatgatccg	catgacgccc
3070	3080	3090	3100	3110	3120
acggtccatcg	ctgtgcattta	ctcggatgaa	acggagcagt	gggagaagtt	cgccctagag
3130	3140	3150	3160	3170	3180
aaqcggcagg	ggcccttgga	gtcatcaag	aaqgggtaca	cccagcagct	ggcccttcaga
3190	3200	3210	3220	3230	3240
caacccagct	ctgcctttgc	ggccttcgtg	aaacggggca	ccaqcacccc	gctgaccggc
3250	3260	3270	3280	3290	3300
tacgtgggca	aggtcttctc	ctcggctgtc	aaactcatcg	ccatcgactc	ccaagtcttc
3310	3320	3330	3340	3350	3360
cgcggggctg	ttaaatggct	gacccctggag	aagcagaagc	ccgacggggt	ctcccgaggag
3370	3380	3390	3400	3410	3420
gatgcccccg	tgatacacca	agaaatgatt	ggtggattac	ggaacaacaa	cgagaaagag
3430	3440	3450	3460	3470	3480
atggccccca	cgccctttct	tctcatctcg	ctgcaggagg	ctaaaatat	ctgcgaggag
3490	3500	3510	3520	3530	3540
caqgtcaaca	gcctggcagg	cagcatcact	aaagcaggag	acttccttga	agccaaactac
3550	3560	3570	3580	3590	3600
atgaacccac	agagatccca	cactgtggcc	attgctggct	atccctctgg	ccagatgggg
3610	3620	3630	3640	3650	3660
aggtccaaag	ggcctcttct	taacaaattt	ctgaccacag	ccaaagataa	gaaccgctgg
3670	3680	3690	3700	3710	3720
gaggaccccc	ctaagcagct	ctacaacgtg	gagggcacat	cctatccccc	cttggcccta
3730	3740	3750	3760	3770	3780
ccccagctaa	aaqactttca	ctttgtggct	ccctctgtgc	gttggctcaa	tgaacagaga
3790	3800	3810	3820	3830	3840
caatcgggtg	gtggctatgg	ctctacccag	gccaccttca	tggtgttcca	agccctggct
3850	3860	3870	3880	3890	3900

6/19

FIG. 2(IV)

caataccaaa	aggacgcccc	tgaccaccag	gaactgaacc	ttgatgtctc	cctccaactg
3910	3920	3930	3940	3950	3960
cccagccgca	gctccaagat	caccacccgt	atccactggg	aactctgccag	cctcctcgga
3970	3980	3990	4000	4010	4020
tcagaagaga	ccaaggaaaa	tgagggtttc	acagtcacag	ctgaaggaaa	aggccaaggc
4030	4040	4050	4060	4070	4080
accttgtcgg	tggtgacaat	gtaccatgct	aaggccaaag	atcaactcac	ctgtaataaa
4090	4100	4110	4120	4130	4140
ttcgacctca	aggtaccat	aaaaccagca	ccggaaacag	aaaagaggcc	tcaggatgcc
4150	4160	4170	4180	4190	4200
aagaacacta	tgatccttga	gatctgtacc	aggtaccggg	gagaccagga	tgccactatg
4210	4220	4230	4240	4250	4260
tctatattgg	acatatccat	gatgactggc	tttctccag	acacagatga	cctgaagcag
4270	4280	4290	4300	4310	4320
ctggccaatg	gtgttgacag	atcacatctc	aagtatgagc	tgacaaaagc	cttctccgat
4330	4340	4350	4360	4370	4380
aggaacaccc	tcacatctta	cctggacaag	gtctcacact	ctgaggatga	ctctctagct
4390	4400	4410	4420	4430	4440
ttcaaaagttc	accaatactt	taatgtagag	cttatccagc	ctggagcagt	caaggtctac
4450	4460	4470	4480	4490	4500
gcctattaca	acctggagga	aagctgtacc	cggttctacc	atccggaaaa	ggaggatgga
4510	4520	4530	4540	4550	4560
aagctgaaca	agctctgccg	tcatgaactg	tgccctgtg	ctgaggagaa	ttgcttcata
4570	4580	4590	4600	4610	4620
caaaaagtcgg	atgacaaggt	cacctggaa	gaacggctgg	acaaggcctg	tgaaccagga
4630	4640	4650	4660	4670	4680
gtggactatg	tgtacaagac	ccgactggtc	aaggttcagc	tgtccaatga	ctttgacgag
4690	4700	4710	4720	4730	4740
tacatcctgg	ccattgagca	gaccatcaag	tcaggctcgg	atgaggtgca	ggtcggacag
4750	4760	4770	4780	4790	4800
cagccacagt	tcacacagcc	catcaagtgc	agagaagccc	tgaagctgga	ggagaagaaa
4810	4820	4830	4840	4850	4860
cactacctca	tgtgggtct	ctcctccgat	ttctggggag	agaagcccaa	cctcagctac
4870	4880	4890	4900	4910	4920
atcctcggga	aggacacttg	ggtggagcac	tggcctgagg	aggacgaatt	ccaagacgaa
4930	4940	4950	4960	4970	4980
gagaaccaga	aacaatgccg	ggacctcggc	gccttcaccc	agagcatggt	tgtctttggg
4990	5000	5010	5020	5030	5040
tcgcccact	gaccacccc	ccattec			
5050	5060				

SUBSTITUTE SHEET (RULE 26)

7/19




FIG. 3

8 / 19




FIG. 4

9 / 19




FIG. 5

10/19




FIG. 6

11/19




FIG. 7

12 / 19




FIG. 8

SUBSTITUTE SHEET (RULE 26)

13 / 19




FIG. 9

SUBSTITUTE SHEET (RULE 26)

14 / 19




FIG. 10

15 / 19




FIG. 11

16 / 19



FIG. 12

17/ 19




FIG. 13

18 / 19




FIG. 14

19/19




FIG. 15

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 97/00603

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C12N15/57 C07K14/47 C12N9/64 A61K38/48
A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF IMMUNOLOGY, vol. 153, 1 December 1994, BALTIMORE US, pages 5285-5302, XP002032794 TANIGUCHI-SIDLE, A. & ISENMAN, D.: "Interactions of human complement component C3 with factor B and with complement receptors type 1 (CR1,CD35) and type 3 (CR3,CD11b/CD38) involve an acidic sequence at the N-terminus of C3 alpha'-chain" cited in the application see the whole document --- -/--	1-4,9, 10,35-38

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

11 June 1997

Date of mailing of the international search report

01.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van der Schaal, C

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/GB 97/00603

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>US 4 661 347 A (MULLER-EBERHARD HANS J ET AL) 28 April 1987</p> <p>see column 3, line 16 - line 24 see column 3, line 63 - column 4, line 20 see column 4, line 50 - line 66 see claims</p> <p style="text-align: center;">---</p>	<p>1-4, 35-41, 44,45,48</p>
X	<p>J IMMUNOL 142 (8). 1989. 2759-2765. CODEN: JOIMA3 ISSN: 0022-1767, XP002032795</p> <p>PANGBURN M K: "ANALYSIS OF THE MECHANISM OF RECOGNITION IN THE COMPLEMENT ALTERNATIVE PATHWAY USING C3B -BOUND LOW MOLECULAR WEIGHT POLYSACCHARIDES."</p> <p>see the whole document</p> <p style="text-align: center;">---</p>	<p>1-4</p>
X	<p>DATABASE BIOSIS</p> <p>BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US</p> <p>FRIES L F ET AL: "COMPLEMENT C-3B COVALENTLY BOUND TO IMMUNOGLOBULIN G DEMONSTRATES A REDUCED RATE OF INACTIVATION BY FACTORS H AND I."</p> <p>XP002032796</p> <p>see abstract</p> <p>& J EXP MED 160 (6). 1984 (RECD. 1985). 1640-1655. CODEN: JEMEAV ISSN: 0022-1007,</p> <p style="text-align: center;">---</p>	<p>1-4</p>
P,X	<p>WO 96 07738 A (IMUTRAN LTD ;HARRISON RICHARD ALEXANDER (GB); FARRIES TIMOTHY CHAR) 14 March 1996</p> <p>see the whole document</p> <p style="text-align: center;">-----</p>	<p>1-12, 35-50</p>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 97/00603

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 49, 50
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 49 and 50 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 97/00603

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4661347 A	28-04-87	NONE	
WO 9607738 A	14-03-96	AU 3477295 A CA 2197888 A	27-03-96 14-03-96